arXiv:math/0007156v1 [math.AG] 26 Jul 2000

Painlevé equations and deformations of rational surfaces with rational double points*†

Masa-Hiko SAITO ‡
Department of Mathematics, Faculty of Science, Kobe University

Hiroshi UMEMURA§
Graduate School of Mathematics, Nagoya University

In this paper we give an answer to the fundamental questions about the Painlevé equations. Where do the Bäcklund transformations come from? Our approach depends on the geometry of the projective surface constructed by Okamoto and reviewed in [U2].

The Painlevé equations were discovered around 1900 in the pursuit of special functions. Painlevé and Gambier classified algebraic differential equations $y'' = R(t, y, y')$ without movable singular points, where R is a rational function of t, y, y' and t is the independent variable. After the refinement procedure of throwing away among the differential equations those that are integrable by the so far known functions, they arrived at the list of the six Painlevé equations P_J $(I \leq J \leq VI)$. After the discovery of Painlevé equations, an unexpected but important feature of the Painlevé equations was revealed. The Painlevé equations have symmetries or the Bäcklund transformations. In fact, if we recall the motivation of the discovery, it is surprising that the Painlevé equations admit symmetries. So a natural question arise. Where do the Bäcklund transformations come from? Our answer is that they arise from the rational double points.

The Painlevé equation P_J is written in a Hamiltonian form

$$\begin{cases} \dfrac{dq}{dt} & = & \dfrac{\partial H_J}{\partial p}, \\[2ex] \dfrac{dp}{dt} & = & -\dfrac{\partial H_J}{\partial q} \end{cases}$$

---

*1991 *Mathematical Subject Classification.* 14D15, 34M55, 32G10, 14J26

†*Key Words and phrases.* Painlevé equations, Bäckglund transformations, Rational Surfaces, Rational double points, Deformations

‡Partly supported by Grant-in Aid for Scientific Research (B-09440015), (B-12440008) and (C-11874008), the Ministry of Education, Science and Culture, Japan

§Partly supported by Grant-in Aid for Scientific Research (B-11440006), the Ministry of Education, Science and Culture, Japan

for an appropriate polynomial H_J of q, p and t ($I \leq J \leq VI$). We also clarify where the polynomial H_J comes from and why it is uniquely determined. This point is a geometric interpretation of [ST] and [MMT]. It is interesting to notice that our theory provides us a lot of pairs (U, V) of algebraic varieties defined over $\mathbb{C}$ such that U and V are not isomorphic as algebraic varieties but they are isomorphic as complex manifolds (cf §11). (The idea in §11 is developed further in the direction of deformation theory of Okamoto–Painlevé pairs and its relation to Painlevé equations in [STT].)

We work with the second Painlevé equation P_{II} to illustrate the general case. Our argument is given in a form easily applicable to the other Painlevé equations (cf. [STT]). Through out the paper, the ground field in $\mathbb{C}$. The most natural setting seems to be, however, over the ring $\mathbb{Z}[\frac{1}{2}]$.

We should mention that a recent work of Sakai [Sakai] is also working on the discrete and differential Painlevé equations related to the geometry of rational surfaces and symmetries of affine Weyl groups.

The authors express gratitude to Professor Kei-ichi Watanabe. The discussions with him was indispensable to write §9.

§1. Construction of a family of rational surfaces.

The second Painlevé equation is an ordinary differential equation

$$P_{II}(\alpha) \qquad y'' = 2y^3 + ty + \alpha$$

of the second order, where t is the independent variable, $y'' = d^2y/dt^2$ and $\alpha \in \mathbb{C}$ is a parameter. The extended affine Weyl group $\widehat{G}$ of type $A_1^{(1)}$ appears as the symmetry of $P_{II}(\alpha)$. Namely if y is a solution of $P_{II}(\alpha)$, then

$$T_+(y) = -y - \frac{\alpha + \frac{1}{2}}{y' + y^2 + \frac{t}{2}}$$

is a solution of $P_{II}(\alpha + 1)$,

$$T_-(y) = -y + \frac{\alpha - \frac{1}{2}}{y' - y^2 - \frac{t}{2}}$$

is a solution of $P_{II}(\alpha - 1)$ and

$$I(y) = -y$$

is a solution of $P_{II}(-\alpha)$. The automorphisms i, t_+, t_- of the affine line $\mathbb{A}^1$ with coordinate system α, i.e. $\mathbb{A}^1 = \text{Spec } \mathbb{C}[\alpha]$, given by

$$t_+(\alpha) = \alpha + 1, \quad t_-(\alpha) = \alpha - 1, \quad i(\alpha) = -\alpha$$

generate the (extended) affine Weyl group $\widehat{G}$ of type $A_1^{(1)}$. So the group $\widehat{G}$ operates on the affine line $\mathbb{A}^1$. We extend the operation of $\widehat{G}$ on the affine plane $\mathbb{A}^2$ with coordinate system (t, α) so that $\widehat{G}$ leaves t invariant. We consider the affine space $\mathbb{A}^4$ with coordinate system (t, α, y, y') and the projection $p_{12} : \mathbb{A}^4 \to \mathbb{A}^2,\ (t, \alpha, y, y') \mapsto (t, \alpha)$. Through the transformations T_+, T_-, I regarded as birational automorphism of the affine space $\mathbb{A}^4$, the extended affine Weyl group $\widehat{G}$ operates birationally on $\mathbb{A}^4$ such that the projection $p_{12} : \mathbb{A}^4 \to \mathbb{A}^2$ is $\widehat{G}$-equivariant. For example the birational automorphism of $\mathbb{A}^4$ induced by T_+ is given by

$$\begin{array}{ccc} \mathbb{A}^4 & \to & \mathbb{A}^4, \\ & & \\ (t, \alpha, y, y') & \mapsto & (t, \alpha+1, T_+(y), T_+(y)') \end{array}$$

where

$$T_+(y) = -y - \frac{\alpha + \frac{1}{2}}{y' + y^2 + \frac{t}{2}}$$

as is given above and

$$\begin{aligned} T_+(y)' &= \left(-y - \frac{\alpha + \frac{1}{2}}{y' + y^2 + \frac{t}{2}}\right)' \\ &= -y' + \frac{(\alpha + \frac{1}{2})(y'' + 2yy' + \frac{1}{2})}{(y' + y^2 + \frac{t}{2})^2} \\ &= -y' + \frac{(\alpha + \frac{1}{2})(2y^3 + ty + \alpha + 2yy' + \frac{1}{2})}{(y' + y^2 + \frac{t}{2})^2}. \end{aligned}$$

Namely if we use a more rigorous notation, the birational automorphism of $\mathbb{A}^4$ is defined by

$$\mathbb{A}^4 \to \mathbb{A}^4, \qquad (t, \alpha + 1, u, v) \mapsto (t, -\alpha, U(t, \alpha, u, v), V(t, \alpha, u, v))$$

where

$$\begin{aligned} U(t, \alpha, u, v) &= -u - \frac{\alpha + \frac{1}{2}}{v + u^2 + \frac{t}{2}}, \\ V(t, \alpha, u, v) &= -v + \frac{(\alpha + \frac{1}{2})(2u^3 + tu + \alpha + 2uv + \frac{1}{2})}{(v + u^2 + \frac{t}{2})^2}. \end{aligned}$$

Remark (1.1). *For the root system of type A_1, the affine Weyl group is isomorphic to the extended affine Weyl group. For a systematic understanding of the Painlevé equations, we should regard $\widehat{G}$ as the extended affine Weyl group.*

On the affine space $\mathbb{A}^4$, we have a vector field

$$\delta(\alpha) = \frac{\partial}{\partial t} + y' \frac{\partial}{\partial y} + (2y^3 + ty + \alpha)\frac{\partial}{\partial y'}$$

so that the birational maps T_+, T_-, I are compatible with the vector field $\delta(\alpha)$. We projectified the fibration $p = p_{12} : \mathbb{A}^4 \to \mathbb{A}^2$ in the following manner ([U2]). To explain the projectification of p, we had better take other coordinate systems on $\mathbb{A}^4$ and $\mathbb{A}^2$. First, we introduce a new coordinate system (t, c) such that $c = \alpha - \frac{1}{2}$. Second, we define a new coordinate system (t, c, p, q) on $\mathbb{A}^4$ such that

$$(1.2) \qquad \begin{cases} q &= y, \\ p &= y' - q^2 - \frac{t}{2}. \end{cases}$$

In terms of this coordinate system, the second Painlevé equation is written as

$$S_2(c) \begin{cases} \dfrac{dq}{dt} &= q^2 + p + \dfrac{t}{2}, \\ \dfrac{dp}{dt} &= -2qp + c \end{cases}$$

and the vector field $\delta(\alpha)$ is given by

$$D(c) = \frac{\partial}{\partial t} + (q^2 + p + \frac{t}{2})\frac{\partial}{\partial q} + (-2qp + c)\frac{\partial}{\partial p}.$$

In other words, if we consider an isomorphism

$$\begin{aligned} \varphi : \mathbb{A}^4 &\longrightarrow \mathbb{A}^4 \\ (t, \alpha, y, y') &\longmapsto \left(t, c, y, y' - y^2 - \frac{t}{2}\right), \end{aligned}$$

then φ transforms $\delta(\alpha)$ to $D(c)$. The system $S_2(c)$ is a Hamiltonian form. In fact if we take $H(t, c, q, p) = q^2 p + \frac{1}{2}p^2 + \frac{t}{2}p - cq$, then we have

$$S_2(c) \begin{cases} \dfrac{dq}{dt} &= \dfrac{\partial H}{\partial p} \\ \dfrac{dp}{dt} &= -\dfrac{\partial H}{\partial q}. \end{cases}$$

We explain now how to projectify the fibration $p_{12} : \mathbb{A}^4 \to \mathbb{A}^2$. We take a point $(t_0, c_0) \in \mathbb{A}^2$ fixed once for all and show how to projectify the fiber $X[t_0, c_0] := p_{12}^{-1}(t_0, c_0)$ of the morphism $p_{12} := \mathbb{A}^4 \to \mathbb{A}^2$, $(t, c, q, p) \mapsto (t, c)$. The fiber $X[t_0, c_0]$ is isomorphic to the affine plane $\mathbb{A}^2$. We need four copies W_i $(1 \le i \le 4)$ of $\mathbb{A}^2$ with coordinate system (y_i, z_i). We glue together the W_i' s by the following rule to get a rational ruled surface $Z[t_0, c_0]$.

(i) A point $(y_1, z_1) \in W_1$ and a point $(y_2, z_2) \in W_2$ are identified if

$$y_1 = y_2 \quad \text{and} \quad z_1 z_2 = 1.$$

(ii) A point $(y_1, z_1) \in W_1$ and a point $(y_3, z_3) \in W_3$ are identified if

$$y_1 y_3 = 1 \quad \text{and} \quad z_1 = c_0 y_3 - y_3^2 z_3.$$

We notice that the latter condition is equivalent to

$$z_3 = c_0 y_1 - y_1^2 z_1.$$

(iii) A point $(y_3, z_3) \in W_3$ and a point $(y_4, z_4) \in W_4$ are identified if

$$y_3 = y_4 \quad \text{and} \quad z_3 z_4 = 1.$$

The projections

$$W_i \to \mathbb{A}^1, \quad (y_i, z_i) \mapsto y_i \quad \text{for } 1 \le i \le 4$$

glue together to give a fibration

$$p_{Z[t_0,c_0]} : Z[t_0, c_0] \to \mathbb{P}^1.$$

So $Z[t_0, c_0]$ is a $\mathbb{P}^1$-bundle over $\mathbb{P}^1$ on $Z[t_0, c_0]$ is a rational ruled surface. We have, as is easily seen

$$Z[t_0, c_0] \cong \begin{cases} F_2 & \text{if } c_0 = 0, \\ \\ \mathbb{P}^1 \times \mathbb{P}^1 & \text{otherwise.} \end{cases}$$

It is apparent that the construction of $Z[t_0, c_0]$ depends only on c_0 and not on t_0. We identify the fiber $X[t_0, c_0]$, which is the affine plane with coordinate system (q, p), with W_1 that is the affine space with coordinate system (y_1, z_1) by sending a point (q, p) of $X[t_0, c_0]$ to the point $(y_1, z_1) = (q, p)$ of W_1. We thus constructed a projectification $Z[t_0, c_0]$ of the fiber $X[t_0, c_0]$ but this is not the desired projectification. To get the projectification $X[t_0, c_0]$, we have to blow-up the rational ruled surface $Z[t_0, c_0]$ eight times, the centers being carefully chosen infinitely near points of $(y_4, z_4) = (0, 0) \in W_4$. The centers depend not only on c_0 but also t_0 so that the projectification $\mathfrak{X}[t_0, c_0]$ depends on c_0 and t_0. The center a_1 of the first blowing-up is the point $(y_4, z_4) = (0, 0)$ on W_4. For $1 \le i \le 8$, we denote by $\pi_i : Z_i[t_0, c_0] \to Z_{i-1}[t_0, c_0]$ the i-the blowing-up, by $E_i \subset Z_i[t_0, c_0]$ the exceptional divisor of π_i, and by $a_i \in Z_{i-1}[t_0, c_0]$ the center of the i-th blown-up π_i, so that $Z_0[t_0, c_0] = Z[t_0, c_0]$, $Z_8[t_0, c_0] = \mathfrak{X}[t_0, c_0]$ and $a_1 = (0, 0) \in W_4$. The curve $S = \{(y_2, z_2) \in W_3 \,|\, z_3 = 0\} \cup \{(y_4, z_4) \,|\, z_4 = 0\}$ is a section of the ruled surface $\pi : Z[t_0, z_0] \to \mathbb{P}^1$ and the self-intersection number $S^2 = 2$. For $2 \leq i \leq 8$ the center $a_i \in Z_{i-1}[t_0, c_0]$ is always on the exceptional divisor E_{i-1} of the previous blowing-up. For $i = 2, 3, 4$, the center $a_i \in Z_{i-1}[t_0, c_0]$ is the intersection point of E_{i-1} and the proper transform of S by the morphism $Z_{i-1}[t_0, c_0] \to Z_0[t_0, c_0] = Z[t_0, c_0]$. Let $D_i \subset \mathfrak{X}[t_0, c_0] = Z_8[t_0, c_0]$ be proper transform of E_i by the morphism $\mathfrak{X}[t_0, c_0] = Z_8[t_0, c_0] \to Z_i[t_0, c_0]$ for $1 \leq i \leq 7$. The following result is proved in [U2].

Sublemma (3.6) **of** [U2]. *Locally on* W_4*, the construction* $\mathfrak{X}[t_0, c_0] = Z_8[t_0, c_0] \to Z[t_0, c_0]$ *or the blowing-up of* $Z[t_0, c_0]$ *is equivalent to the minimal resolution of the rational map*

$$F : W_4 \cdots \to \mathbb{P}^1, \quad (y_4, z_4) \mapsto (y_4^4 z_4, 2z_4 - y_4^4 + t y_4^2 z_4 + (2c_0 + 1) y_4^3 z_4).$$

Remark. *F is not regular on $\mathfrak{X}[t_0, c_0]$. For, it has a base point $(y_2, z_2) = 0$ on W_2. This point outside W_4 is left untouched in the construction of $\mathfrak{X}[t_0, c_0]$.*

We denote by D_0 the proper transform of the curve $S \subset Z[t_0, c_0]$ by the morphism $\mathfrak{X}[t_0, c_0] \to Z[t_0, c_0]$. Then we have $D_1 \simeq \mathbb{P}^1$ and $D_i^2 = -2$ for $1 \leq i \leq 7$. The configuration of the D_i's is shown in Fig.(1.3).



Fig.(1.3)

The dual graph of Fig.(1.3) is Fig.(1.4) that is the Dynkin diagram of type $E_7^{(1)}$.

Namely the vertices in Fig.(1.4) correspond to the curves in Fig.(1.3). Two different vertices are jointed by an edge if corresponding two curves have a point in common. We set $\mathfrak{D}[t_0, c_0] := \cup_{i=0}^7 D_i \subset \mathfrak{X}[t_0, c_0]$, which is a divisor on $\mathfrak{X}[t_0, c_0]$. The divisors D_i also depend on t_0 and c_0. So we should denoted D_i by $\mathfrak{D}_i[t_0, c_0]$. To



Fig.(1.4)

simplify the notation, we use $\mathfrak{D}_i[t_0, c_0]$, only when we emphasize the dependence on t_0 and c_0. So far we fixed $(t, c) = (t_0, c_0)$. The above construction works globally on the fibration $\pi : \mathbb{A}^4 \to \mathbb{A}^2$ and gives us a fiber space

$$\varphi : \mathfrak{X} \to \mathbb{A}^2$$

and a divisor $\mathfrak{D}$ on $\mathfrak{X}$ such that for a point $(t_0, c_0) \in \mathbb{A}^2$, the fiber $\varphi^{-1}(t_0, c_0)$ is isomorphic to $\mathfrak{X}[t_0, c_0]$ and such that $\mathfrak{D} \cap \varphi^{-1}(t_0, c_0)$ yields the divisor $\mathfrak{D}[t_0, c_0]$ on $\mathfrak{X}[t_0, c_0]$. Namely, let ψ be the composite $p_2 \circ \varphi$

$$\mathfrak{X} \xrightarrow{\varphi} \mathbb{A}^2 \xrightarrow{p_2} \mathbb{A},$$

where $p_2 : \mathbb{A}^2 \to \mathbb{A}^1$ is the projection onto the second factor so that $p_2(t, c) = c$ for a point $(t, c) \in \mathbb{A}^2$. We denote the fiber $\psi^{-1}(c_0)$ by $\mathfrak{X}[c_0]$ for a point $c_0 \in \mathbb{A}^1$. Then φ gives a morphism

$$\varphi_{c_0} : \mathfrak{X}[c_0] \to \mathbb{A}^1 \times c_0 \subset \mathbb{A}^2.$$

Since $\mathbb{A}^1 \times c_0 \simeq$ Spec $\mathbb{A}[t]$ is the affine line with coordinate system t, we have a morphism

$$\varphi[c_0] : \mathfrak{X}[c_0] \to \text{Spec } \mathbb{C}[t],$$

which we denote also by φ_{c_0}. On the threefold $\mathfrak{X}[c_0]$, we have a divisor $\mathfrak{D}[c_0] := \mathfrak{D} \cap \mathfrak{X}[c_0]$ and a rational vector field

$$\delta[c_0] = \frac{\partial}{\partial t} + (y_1^2 + z_1 + \frac{t}{2})\frac{\partial}{\partial y_1} + (-2y_1z_1 + c_0)\frac{\partial}{\partial z_1}$$

generates a foliation on $\mathfrak{X}[c_0]$. The divisor $\mathfrak{D}[c_0]$ is the set of singular points of the foliation so that every leaf is transversal to the fibers of $\mathfrak{X} \to$ Spec $\mathbb{C}[t]$ in the open set $\mathfrak{X}[c_0] \backslash \mathfrak{D}[c_0]$. Since the Painlevé equations have no movable singular points (=movable branch points and movable essential singular points), the set of solutions on the set of leaves of the foliation in $\mathfrak{X}[c_0] \backslash \mathfrak{D}[c_0]$ sweeps the whole space $\mathfrak{X}[c_0] \backslash \mathfrak{D}[c_0]$ (See Fig.(1.5)).



Fig.(1.5)

$\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ is the space of initial conditions of the second Painlevé equation introduced by Okamoto [O] (cf. [U2]). In terms of coordinate system (t, c, q, p) of the affine space $\mathbb{A}^4$, the affine extended Weyl group G is generated by rational maps

$$J : \mathbb{A}^4 \cdots \to \mathbb{A}^4, \quad (t, c, q, p) \mapsto (t, -1 - c, -q, -2q^2 - p - t),$$

and

$$I : \mathbb{A}^4 \cdots \to \mathbb{A}^4, \quad (t, c, q, p) \mapsto (t,\ -c,\ q - \frac{c}{p},\ p).$$

The corresponding birational map induced respectively by J (resp. I) on a variety $\mathfrak{Z}$ over $\mathbb{A}^2$, which is $\mathbb{A}^2$-birational to $p_{12} : \mathbb{A}^4 \to \mathbb{A}^2$, will be denoted by $J_{\mathfrak{Z}}(c, -1 - c)$ (resp. $I_{\mathfrak{Z}}(c, -c)$). In this context, when we specialize the parameter c to $c_0 \in \mathbb{C}$, we denote the thus obtained birational map by $J_{\mathfrak{Z}}(c_0, -1 - c_0)$, $I_{\mathfrak{Z}}(c_0, -c_0)$. We proved in [U2] that when the parameter c takes a fixed value $c_0 \in \mathbb{C}$, the Bäcklund transformations give the following isomorphisms that commute with the foliations.

$$J_{\mathfrak{X}}(c_0, -1 - c_0) : \mathfrak{X}[c_0] \to \mathfrak{X}[-1 - c_0]. \tag{1.6}$$

$$I_{\mathfrak{X}}(c_0, -c_0) : \mathfrak{X}[c_0] \to \mathfrak{X}[-c_0]. \tag{1.7}$$

Moreover we have

$$J_{\mathfrak{X}}(c_0, -1 - c_0)[\mathfrak{D}[c_0]] = \mathfrak{D}[-1 - c_0] \tag{1.8}$$

and

$$(1.9) \qquad I_{\mathfrak{X}}(c_0, -c_0)[\mathfrak{D}[c_0]] = \mathfrak{D}[-c_0].$$

For, for every $c_0 \in \mathbb{C}$, $\mathfrak{D}[c_0]$ is the set of singular points of the foliation on $\mathfrak{X}[c_0]$ and the isomorphisms $J_{\mathfrak{X}}(c_0, -1-c_0)$ and $I_{\mathfrak{X}}(c_0, -c_0)$ commute with the foliation.

Proposition (1.10). *We have*

$$\begin{aligned} J_{\mathfrak{X}}(c_0, -1-c_0)(\mathfrak{D}_i[c_0]) &= \mathfrak{D}_{8-i}[-1-c_0] \quad \textit{for } 1 \le i \le 7, \\ J_{\mathfrak{X}}(c_0, -1-c_0)(\mathfrak{D}_0[c_0]) &= \mathfrak{D}_0[-1-c_0], \\ I_{\mathfrak{X}}(c_0, -c_0)(\mathfrak{D}_i[c_0]) &= \mathfrak{D}_i[-c_0] \quad \textit{for } 0 \le i \le 7. \end{aligned}$$

Proof. Once we have (1.8) and (1.9), we can check the identities easily.

When we specialize not only c to c_0 but also t to t_0, the corresponding birational maps are denoted respectively by $J_3(t_0; c_0, -1-c_0)$ and $I_3(t_0; c_0, -c_0)$. Using this notation, we proved in [U2] that we have isomorphisms

$$J_{\mathfrak{X}}(t_0; c_0, -1-c_0) : \mathfrak{X}[t_0, c_0] \to \mathfrak{X}[t_0, -1-c_0]$$

and

$$I_{\mathfrak{X}}(t_0; c_0, -c_0) : \mathfrak{X}[t_0, c_0] \to \mathfrak{X}[t_0, -c_0].$$

The following results follows from Proposition (1.10).

Proposition (1.11). *When t and c take fixed values t_0 and c_0, we have the following identities.*

(i) $J_{\mathfrak{X}}(t_0; c_0, -1-c_0)(\mathfrak{D}_i[t_0, c_0]) = \mathfrak{D}_{8-i}[t_0, -1-c_0]$

for $1 \le i \le 7$ and

(ii) $J_{\mathfrak{X}}(t_0; c_0, -1-c_0)(\mathfrak{D}_0[t_0, c_0]) = \mathfrak{D}_0[t_0, -1-c_0]$.

(iii) $I_{\mathfrak{X}}(t_0; c_0, -c_0)(\mathfrak{D}_i)[t_0, -c_0]$.

Remark (1.12). *$J_{\mathfrak{X}}; \mathfrak{X} \to \mathfrak{X}$ is a regular automorphism and the morphism $\varphi : \mathfrak{X} \to \mathbb{A}^2$ is compatible with automorphism $j : \mathbb{A}^2 \to \mathbb{A}^2$, $(t, c) \mapsto (t, -c)$. On the other hand, $I_{\mathfrak{X}} : \mathfrak{X} \to \mathfrak{X}$ is not biregular.*

In fact there exists a curve $C_2'[t_0, 0]$ on $\mathfrak{X}[t_0, 0]\backslash\mathfrak{D}$ isomorphic to $\mathbb{P}^1$ giving the Riccati solutions of $P_{II}\left(\frac{1}{2}\right)$. The curve $C_2'[t_0, 0]$ is the base locus of the rational map $I_{\mathfrak{X}} : \mathfrak{X} \to \mathfrak{X}$. We proved in [U2] that if we denote by $\widetilde{\mathfrak{X}}$ the blowing-up of $\mathfrak{X}$ along the surface $\bigcup_{t_0\in\mathbb{C}} C_2'[t_0, 0]$, then $I_{\widetilde{\mathfrak{X}}} : \widetilde{\mathfrak{X}} \to \widetilde{\mathfrak{X}}$ is a biregular automorphism. We explain §3 the reason why we have to blow up $\mathfrak{X}$ to make $I_{\mathfrak{X}}$ biregular. The birational maps

$$I_{\mathfrak{X}}(c_0, -c_0) : \mathfrak{X}[c_0] \to \mathfrak{X}[-c_0]$$

and

$$I_{\mathfrak{X}}(t; c_0, -c_0) : \mathfrak{X}[t_0, c_0] \to \mathfrak{X}[t_0, c_0]$$

induced by $I_{\mathfrak{X}}$ is the identity, when $c_0 = 0$. So they are biregular. Hence strictly speaking in the proofs of Propositions (1.10), (1.11) we must treat the case $c_0 = 0$ separately. For every integer n, there is curve $C[t_0, n]$ isomorphic to $\mathbb{P}^1$ on $\mathfrak{X}[t_0, n]\backslash\mathfrak{D}$. We blow up $\mathfrak{X}$ along the surfaces $\bigcup_{t_0\in\mathbb{C}} C[t_0, n]$ for every $n \in \mathbf{Z}$. We get a manifold $\mathcal{Y}$, a projective limit of a scheme over $\mathbb{A}^2$ such that the extended Weyl group $\widehat{G} =< I_{\mathcal{Y}}, J_{\mathcal{Y}} >$ operates regularly on $\mathcal{Y}$. In fact, we have seen in [U2], $I_{\mathcal{Y}}$ and $J_{\mathcal{Y}}$ are automorphisms of $\mathcal{Y}$. The extended affine Weyl group $\widehat{G}$ operates on $\mathcal{Y}$ and $\mathbb{A}^2$ in such a way that $\varphi_{\mathcal{Y}} : \mathcal{Y} \to \mathbb{A}^2$ is $\widehat{G}$-equivariant. Here $\varphi_{\mathcal{Y}} : \mathcal{Y} \to \mathbb{A}^2$ is the composite of the blowing-up morphism $\mathcal{Y} \to \mathfrak{X}$ and the morphism $\varphi : \mathfrak{X} \to \mathbb{A}^2$ arising from the projection $p_{12} : \mathbb{A}^4 \to \mathbb{A}^2$. It is natural to ask a

Question (1.13). *The quotient spaces $\mathcal{Y}/\widehat{G}$, $\mathbb{A}^2/\widehat{G}$ and the quotient morphism $\mathcal{Y}/\widehat{G} \to \mathbb{A}^2/\widehat{G}$ are algebraizable?*

§2. Affine root systems on $\mathfrak{X}[t_0, c_0]$.

The additive group Pic $\mathfrak{X}[t_0, c_0]$ of linear equivalence classes of divisors on the surface $\mathfrak{X}[t_0, c_0]$ is a lattice, of rank 10, the bilinear form on Pic $\mathfrak{X}[t_0, c_0]$ being defined by the intersection pairing.

Lemma (2.1). *We can blow down the surface $\mathfrak{X}[t_0, c_0]$ to $\mathbb{P}^2$.*

Proof. If $c_0 \neq 0$, then the ruled surface $Z[t_0, c_0]$ is $\mathbb{P}^1 \times \mathbb{P}^1$. Let $\widetilde{X}$ be the blown-up of $\mathbb{P}^1 \times \mathbb{P}^1$ at a point $P = (x_1, x_2)$ and let C_1, C_2 be respectively the proper transform of $x_1 \times \mathbb{P}^1$, $\mathbb{P}^1 \times x_2$ under the blowing down morphism $\varphi : \widetilde{X} \to \mathbb{P}^1 \times \mathbb{P}^1$. Then C_1 and C_2 are disjoint -1-curves so that we can collapse C_1 and C_2 on $\widetilde{X}$ to get $\mathbb{P}^2$. If the parameter $c_0 = 0$, then the ruled surface $Z[t_0, c_0]$ is isomorphic to F_2. We blow-up F_2 at a point P on the section S of $\pi : F_2 \to \mathbb{P}^1$ with $S^2 = 2$ to get a surface $\widetilde{X}$. So we have the blowing-down morphism $\varphi : \widetilde{X} \to F$. Since $F_2\backslash S = Z[t_0, c_0]\backslash S$ is a line bundle over $\mathbb{P}^1$, we can find a section S_0 of $\pi : F_2 \to \mathbb{P}^1$ such that $S_0^2 = -2$and $(S.S_0) = 0$. Let C_1 be the proper transform of the fiber of the ruled surface $Z[t_0, c_0]$ passing through the point P and let C_2 be the proper transform of S_0 under $\varphi : \widetilde{X} \to F_2$. Then since $C_1^2 = -2$, $C_2^2 = -2$ and $(C_1.C_2) = 1$, we can collapse successively C_1 and C_2 to get $\mathbb{P}^2$. The surface $\mathfrak{X}[t_0, c_0]$ is obtained by blowing-up the ruled surface $Z[t_0, c_0]$ 8 times. the first blowing-up is one of the blowing-ups described above according as $c_0 \neq 0$ or $c_0 = 0$. Hence we can blow-down the surface $\mathfrak{X}[t_0, c_0]$ to $\mathbb{P}^2$.

Corollary (2.2). *The Picard lattice* Pic $\mathfrak{X}[t_0, c_0]$ *is isomorphic to the lattice* $\bigoplus_{i=1}^{10} \mathbf{Z}e_i$ *of rank 10 such that*

$$(e_i, e_j) = \begin{cases} \delta_{ij} & \text{if } i = 1, \\ -\delta_{ij} & \text{otherwise.} \end{cases}$$

Proof. This is an immediate consequence of Lemma (2.1).

We have seen in §1 that on the rational surface $\mathfrak{X}[t_0, c_0]$ there exists the divisor $\mathfrak{D}[t_0, c_0]$ of which irreducible components $\mathfrak{D}_i[t_0, c_0]$, $0 \le i \le 7$ are isomorphic to $\mathbb{P}^1$ with self-intersection number -2 and their dual graph is $E_7^{(1)}$ (cf. Fig.(1.4)). The linear equivalence classes $[D_i]$'s are linearly independent in the Picard lattice $L =$ Pic $\mathfrak{X}[t_0, c_0]$. So the Picard lattice L contains a free $\mathbf{Z}$-module of rank 8 spanned by the class $[D_i]$ of D_i in Pic $\mathfrak{X}[t_0, c_0]$ $(0 \le i \le 7)$, of which the bilinear form is given by minus of the Cartan matrix of the affine root system of type $E_7^{(1)}$. We denote by abuse notation, this subgroup of L with the induced bilinear form by $L(E_7^{(1)})$. For a subset M of L we set

$$M^{\perp} = \{x \in L \,|\, (x, y) = 0 \quad \text{for every } x \in M\}\,.$$

Our aim is to prove Theorem (2.24) and (2.31). We introduced the curves D_i, $0 \le i \le 7$ on the algebraic surface $\mathfrak{X}[t_0, c_0]$. To prove Theorem (2.24), we need some more curves on $\mathfrak{X}[t_0, c_0]$. On the ruled surface $\pi : Z_0[t_0, c_0] \to \mathbb{P}^1$, we have the following curves. All the irreducible components of the curves are isomorphic to $\mathbb{P}^1$.

$$\text{(2.3.1)} \qquad C_1^0[t_0, c_0] : \; = \; \overline{\{(y_3, z_3) \in W \,|\, y_3 = 0\}} \subset Z_0[t_0, c_0]$$

$$= \; \{(y_3, z_3) \in W_3 \,|\, y_3 = 0\} \cup \{(y_4, z_4) \in W_4 \,|\, y_4 = 0\}\,,$$

which is a fiber of π. Here $\bar{A}$ denotes the Zariski closure of a subset A of $Z[t_0, c_0]$.

$$\text{(2.3.2)} \qquad C_2^0[t_0, c_0] := \overline{\{(y_3, z_3) \in W \,|\, y_3 z_3 - c_0 = 0\}} \subset Z_0[t_0, c_0].$$

so that

$$\text{(2.3.3)} \quad C_2^0[t_0, c_0] \; = \; \{(y_3, z_3) \in W \,|\, y_3 z_3 - c_0 = 0\} \cup \{(y_1, z_1) \in W_1 \,|\, z_1 = 0\}$$

$$\cup \{(y_4, z_4) \in W_4 \,|\, y_4 - c_0 z_4 = 0\}$$

on $Z_0[t_0, c_0]$. The curve $C_2^0[t_0, c_0]$ is isomorphic to $\mathbb{P}^1$ for $c_0 \neq 0$, and the union of 2 curves isomorphic to $\mathbb{P}^1$ for $c_0 = 0$. Namely we have

$$(2.3.4) \qquad C_2^0[t_0, c_0] = \begin{cases} \overline{\{(y_1, z_1) \in W_1 \mid z_1 = 0\}}, & \text{if } c_0 \neq 0, \\ \\ \overline{\{(y_1, z_1) \in W_1 \mid z_1 = 0\}} \cup C_1^0[t_0, c_0], & \text{if } c_0 = 0, \end{cases}$$

on the surface $Z[t_0, c_0]$.

$$(2.3.5) \qquad C_4^0[t_0, c_0] := \overline{\{(y_1, z_1) \in W_1 \mid 2y_1^2 + z_1 + t_0 = 0\}} \subset Z_0[t_0, c_0],$$

$$(2.3.6) \quad C_4^7[t_0, c_0] : \; = \; \text{the proper transform of } C_4^0[t_0, c_0] \text{ under the birational}$$
$$\text{morphism } Z_7[t_0, c_0] \to Z_0[t_0, c_0].$$

$$(2.3.7) \qquad C_5^0[t_0, c_0] := \overline{\{(y_1, z_1) \in W_1 \mid y_1 z_1 - c_0 = 0\}} \subset Z[t_0, c_0].$$

So

$$\begin{aligned} C_5^0[t_0, c_0] \;\; = \;\; & \{(y_1, z_1) \in W_1 \mid y_1 z_1 - c_0 = 0\} \cup \{(y_3, z_3) \in W_3 \mid z_3 = 0\} \\ & \cup \{(y_2, z_2) \in W_2 \mid y_2 - c_0 z_2 = 0\}, \end{aligned}$$

which is isomorphic to $\mathbb{P}^1$ if $c_0 \neq 0$ and, which is the union of 2 curves isomorphic to $\mathbb{P}^1$ if $c_0 = 0$.

$$C_5^0 = \overline{\{(y_1, z_1) \in W_1 \mid y_1 = 0\}} \cup \overline{\{(y_1, z_1) \in W_1 \mid z_1 = 0\}}.$$

$$(2.3.8) \qquad C_6^0[t_0, c_0] := \overline{\{(y_1, z_1) \in W_1 \mid 2y_1^3 + t_0 y_1 + y_1 z_1 + c_0 + 1 = 0\}}.$$

$C_6^0[t_0, -1]$ is irreducible if $c_0 \neq -1$ and if $c_0 = -1$, it has 2 irreducible components:

$$C_6^0[t_0, -1] = \overline{\{(y_1, z_1) \in W_1 \mid y_1 = 0\}} \cup C_4^0[t_0, -1].$$

We denote by $C_3[t_0, c_0]$ the exceptional divisor E_8 of the last or the eighth blow-up $\mathfrak{X}[t_0, c_0] = Z_8[t_0, c_0] \to Z_7[t_0, c_0]$ in the construction of $\mathfrak{X}[t_0, c_0]$. We denote the proper transform of $C_j^{(0)}[t_0, c_0]$ by $C_j[t_0, c_0]$ for $1 \leq j \leq 6$ with $j \neq 3, 4$ under the map $\mathfrak{X}[c_0, t_0] \to Z_0[t_0, c_0]$. $C_4[t_0, c_0]$ is the total transform of $C_4^{(7)}[t_0, c_0]$ under the birational morphism $\mathfrak{X}[t_0, c_0] = Z_8[t_0, c_0] \to Z_7[t_0, c_0]$. To simplify the notation, the curves $C_j^0[t_0, c_0]$, $C_j[t_0, c_0]$ are respectively denoted simply by C_j^0, C_j if there is no danger of confusion. The surface $Z[t_0, c_0] \to \mathbb{P}^1$ is a ruled surface so that Pic $Z[t_0, c_0] = \mathbf{Z}C_1^0 \oplus \mathbf{Z}S$. We have $(C_1^0)^2 = -1$, $S^2 = 2$, $(C_1^0.S) = 1$. We recall that S is section of the ruled surface $Z[t_0, c_0]$ introduced in §1.

Lemma (2.4). *The curve $C_6^0 \cap W_4$ on W_4 is defined by*

$$2z_4 - y_4^4 + ty_4^2 z_4 + (2c_0+1)y_4^3 z_4 = 0.$$

This is the curve appeared in Sublemma (3.6) of [U2].

Proof. This follows from the definition of the curve C_6^0 and the coordinate transformations among W_i $(1 \le i \le 4)$.

Lemma (2.5). *We have $C_2^0 \sim S - C_1^0$ on $Z[t_0, c_0]$.*

Proof. Since Pic $Z[t_0, c_0] \simeq \mathbf{Z}C_1^0 + \mathbf{Z}S$, we can find integers a, b such that $C_2^0 \sim aS + bC_1^0$. Then $1 = (S.C_2^0) = (S.aS + bC_1^0) = aS^2 + b(S.C_1^0) = 2a + b$, $1 = (C_1^0.C_2^0) = (C_1^0.aS + bC_1^0) = a(C_1^0.S) = 1$. Hence $a = 1$, $b = -1$.

Lemma (2.6). *Assume $c_0 \neq 0$ so that C_2^0 is isomorphic to $\mathbb{P}^1$. Let C_1^1 and C_2^1 be the proper transform of C_1^0 and C_2^0 respectively under the birational morphism $\pi_1 : Z_1[t_0, c_0] \to Z_0[t_0, c_0]$.*

(i) *C_1^1, C_2^1 are isomorphic to $\mathbb{P}^1$.*

(ii) *$(C_1^1.C_2^1) = 0$.*

(iii) *${C_1^1}^2 = {C_2^1}^2 = -1$.*

(iv) *$(C_1^1.E_1) = (C_2^1.E_1) = 1$.*

(v) *$C_1^1 \cap E_1$, $C_2^1 \cap E_1$ and the center $a_2 \in E_1$ are three distinct points on E_1.*

Proof. **(i)** follows the fact that the curves C_1^0, C_1^0 are isomorphic to $\mathbb{P}^1$. The center a_1 of the first blowing-up is the point $(y_4, z_4) = (0, 0)$ on W_4. So that C_1^0, C_2^0 pass through a_1. We have $(C_1^0)^2 = 0$ and $(C_2^0)^2 = 0$ by Lemma (2.5). So that $(C_1^1)^2 = (C_2^1)^2 = -1$. Moreover C_1^0 and C_2^0 intersect at a_1 transversely $(C_1^1.C_2^1)) = 0$. Easy calculation shows that C_1^1 and C_1^2 do not pass through $a_2 \in E_1 \cap C_6^1$. Here C_6^1 is the proper transform of C_6^0 under the birational map $\pi_1 : Z_1[t_0, c_0] \to Z_0[t_0, c_0]$ (cf. Lemma (2.4) and Sublemma (3.6) of [U2]).

Corollary (2.7). *If $c_0 \neq 0$, then the configuration of the curves D_i $(0 \le i \le 7)$ and C_1, C_2 is as in Fig.(2.7.1). We have $(C_1)^2 = (C_2)^2 = -1$.*

Proof. In fact, the centers a_i for $2 \le i \le 8$ are infinitely near points of a_2. So C_1, C_2 are total transform of C_1^1 and C_2^1 respectively under the morphism $Z_8[t_0, c_0] \to Z_2[t_0, c_0]$, which is an isomorphism on a neighborhood of $C_1^1 \cup C_2^1$.

Similarly we can prove the following

Lemma (2.8). *If $c_0 = 0$, then $C_2 = C_1 \cup C_2'$ such that C_1,C_2' are isomorphic to $\mathbb{P}^1$ and we have $(C_1.C_2') = 1$, $(C_1)^2 = -1$, $(C_2')^2 = -2$. The configuration of C_1, C_2' is as in Fig.(2.8.1).*



Fig.(2.7.1)



Fig.(2.8.1)

Proof. If $c_0 = 0$, then a point $(y_1, z_1) \in W_1$ and a point $(y_3, z_3) \in W_3$ are identified if $y_1y_3 = 1$ and $z_3 = -y_1z_1$ so that $Z[t_0, 0] \simeq F_2$ and we have 2 disjoint sections of the ruled surface $Z[t_0, 0]$. Namely

$$S = \{(y_2, z_2) \in W_2 \mid z_2 = 0\} \cup \{(y_4, z_4) \in W_4 \mid z_4 = 0\}$$

and

$$S_0 = \{(y_1, z_1) \in W_1 \mid z_1 = 0\} \cup \{(y_3, z_3) \in W_3 \mid z_3 = 0\}\,.$$

Then $S^2 = 2$, $S_0^2 = -2$. By (2.3.4), we have $C_2^0 = S_0 \cup C_1^0$. C'_2 is the total transform of S_0 that coincides with its proper transform because S_0 does not pass through the center a_1 of the first blowing-up and consequently any centers a_j, $2 \leq j \leq 8$. In particular $C'_2 \subset \mathfrak{X} \backslash \left(\bigcup_{i=0}^{7} D_i \right)$ (cf. Proposition (4.2) below).

Let us now study the curve C_4.

Lemma (2.9). *The curve $C_4^0 \cup W_4$ on the affine plane $W_4 \subset Z_0[t_0, c_0]$ is defined by*

$$2z_4 + (c_0 z_4 - y_4)y_4^3 + t y_4^2 z_4 = 0.$$

Proof. Since $C_4^0 = \overline{\{(y_1, z_1) \in W_1 \,|\, 2y_1^2 + z_1 + t = 0\}}$, this follows from the coordinate transformations among the W_i's $1 \leq i \leq 4$ given §1.

Lemma (2.10). *The curve* C_4^0 *has the following properties.*

(i) *The proper transform* C_4^i *of* C_4^0 *on* $Z_i[t_0, c_0]$ *by the birational morphism* $Z_i[t_0, c_0] \to Z_0[t_0, c_0]$ *passes through the center* a_{i+1} *of the* $(i+1)$*-th blow-up for* $0 \leq i \leq 6$.

(ii) *If* $c_0 \neq -1$, C_4^7 *does not pass through* a_8.

(iii) *If* $c_0 = -1$, C_4^7 *pass through* a_8.

Proof. This is a consequence of Lemma (2.9) and Sublemma (3.6) of [U2].

Corollary (2.11). (i) *On* $Z[t_0, c_0]$, *we have* $C_4^0 \sim S + 2C_1^0$ *so that* $\left(C_4^0\right)^2 = 6$.

(ii) $\left(C_4^7\right)^2 = -1$.

(iii) $\left(C_4\right)^2 = -1$.

(iv) $(C_3 . C_4) = 0$.

Proof. Let $C_4^0 \sim aS + bC_1^0$ with $a, b \in \mathbf{Z}$, on $Z_0[t_0, c_0]$. Since for a given y_1, the equation $2y_1^2 + z_1 + t = 0$ for z_1 has a unique solution,

$$1 = \left(C_1^0 \,.\, C_4^0\right) = \left(C_1^0 \,.\, aS + bC_1^0\right) = a.$$

The defining equation for C_4^0 on W_2 is $2y_2^2 z_2 + 1 + t z_2 = 0$. So $C_4^0 \cap W_2 \cap S = \emptyset$. The defining equation of C_4^0 on W_4 is

$$2z_4 + (c_0 z_4 - y_4)y_4^3 + t y_4^2 z_4 = 0$$

so that on W_4, $W_4 \cap C_4^0 \cap S = (0, 0) \in W_4$. To multiplicity of $W_4 \cap C_4^0 \cap S$ is 4. Hence

$$4 = \left(S \,.\, C_4^0\right) = \left(S \,.\, aS + bC_1^0\right) = 2a + b$$

so that $b = 2$. Consequently $C_4^0 \sim S + 2C_1^0$.

$$\left(C_4^0\right)^2 = \left(S + 2C_1^0\right)^2 = S^2 + 4\left(S \,.\, C_1^0\right) + 4\left(C_1^0\right)^2 = 2 + 4 = 6.$$

Therefore $\left(C_4^7\right)^2 = -1$ by Lemma (2.10). (iii) and (iv) follow from (ii) and Lemma (2.10).

Corollary (2.12). *If* $c_0 \neq -1$, **(i)** C_4 *is an irreducible curve,* **(ii)** *we have* $\left(C_3\right)^2 = \left(C_4\right)^2 = -1$ *and* **(iii)** *the configuration of the curves* D_i $(0 \leq i \leq 7)$ *and* C_3, C_4 *is as Fig.(2.12.1).*



Fig.(2.12.1)

Proof. This is a direct consequence of Lemma (2.10) and Corollary (2.11).

Corollary (2.13). *If $c_0 = -1$, then $C_4 = C_3 \cup C_4'$, where C_4' is isomorphic to $\mathbb{P}^1$. We have $(C_3)^2 = -1$, $(C_4')^2 = -2$, $(C_3 \,.\, C_4') = 1$. We also have $(C_3)^2 = (C_4)^2 = -1$. C_4' coincides with a component of C_6*

Proof. The decomposition of C_4 follows from Lemma (2.10). Since C_3 is the exceptional curve of the first kind on $\mathfrak{X}[t_0, c_0]$, $(C_3)^2 = -1$. We can check $(C_3 \,.\, C_4') = 1$ by a direct calculation. By Corollary (2.11), (ii),

$$-1 = (C_4)^2 = (C_3 + C_4')^2 = (C_3)^2 + 2\,(C_3 \,.\, C_4') + (C_4')^2 = -1 + 2 + (C_4')^2 = 1 + (C_4')^2 .$$

So $(C_4')^2 = -2$. The last assertion follows from the definition of C_4' and C_6.

Corollary (2.14). *If $c_0 = -1$, the configuration of the curves D_i $(0 \leq i \leq 7)$ and C_3, C_4' is as in Fig.(2.14.1).*



Fig.(2.14.1)

Lemma (2.15). *We have* **(i)** $(C_1 \,.\, C_3) = 0$, **(ii)** $(C_1 \,.\, C_4) = 0$, **(iii)** $(C_2 \,.\, C_3) = 0$ *and* **(iv)** $(C_2 \,.\, C_4) = 2$.

Proof. By Lemma (2.5) and Corollary (2.11)

$$C_2^0 \sim S - C_1^0, \qquad C_4^0 \sim S + 2C_1^0$$

so that

$$\left(C_2^0 \,.\, C_4^0\right) = \left(S - C_1^0 \,.\, S + 2C_1^0\right) = S^2 + (C_1^0 \,.\, S) = 2 + 1 = 3.$$

Both C_2^0 and C_4^0 pass through the point $(y_4, z_4) = (0, 0)$ on W_4, which is not a common point of C_2^1 and C_4^1. Here C_i^1 is the proper transform of C_i^0 for birational morphism $\pi_1 : Z_1[t_0, c_0] \to Z_0[t_0, c_0]$. Hence $2 = (C_2^1 \,.\, C_4^1) = (C_2 \,.\, C_4)$. Other intersection numbers are evident from the geometric definition of the curves.

As we imagine from Fig.(2.7.1) and (2.12.1), (2.8.1) and (2.14.1). The pairs $\{C_1, C_2\}$, $\{C_3, C_4\}$ of curves are symmetric with respect to $J_{\mathfrak{X}}(t_0; c_0, -1-c_0)$ (cf. Proposition (1.10)).

Proposition (2.16). **(i)** *We have*

$$J_{\mathfrak{X}}(t_0; c_0, -1-c_0)\,(C_1[t_0, c_0]) = C_3[t_0, -1-c_0],$$

$$J_{\mathfrak{X}}(t_0; c_0, -1-c_0)\,(C_2[t_0, c_0]) = C_4[t_0, -1-c_0],$$

$$J_{\mathfrak{X}}(t_0; c_0, -1-c_0)\,(C_3[t_0, c_0]) = C_1[t_0, -1-c_0],$$

$$J_{\mathfrak{X}}(t_0; c_0, -1-c_0)\,(C_4[t_0, c_0]) = C_2[t_0, -1-c_0].$$

(ii) *Moreover we have*

$$J_{\mathfrak{X}[t_0,c_0]}(c_0, -1-c_0)\,(C_5[t_0, c_0]) = C_6[t_0, -1-c_0],$$

$$J_{\mathfrak{X}[t_0,c_0]}(c_0, -1-c_0)\,(C_6[t_0, c_0]) = C_5[t_0, -1-c_0].$$

Proof. These formulas follow from the definition of curves and from the proof of Lemma (3.5) in [U2], where the local forms $J_{ij} : W_i \to W_j$ of $J_{\mathfrak{X}}[t_0; c_0, -1-c_0]$ are given.

We can prove the following proposition in a similar way.

Proposition (2.17). **(i)** *If* $c_0 \neq 0$, *we have*

$$I_{\mathfrak{X}}[t_0; c_0, -c_0]\,(C_2[t_0, c_0]) \;=\; C_1[t_0, -c_0],$$

$$I_{\mathfrak{X}}[t_0; c_0, -c_0]\,(C_1[t_0, c_0]) \;=\; C_2[t_0, -c_0].$$

(ii) *We also have*

$$I_{\mathfrak{X}}[t_0; c_0, -c_0]\,(C_3[t_0, c_0]) = C_3[t_0, -c_0].$$

Remark (2.18). *If* $c_0 = 0$, $I_{\mathfrak{X}}[t_0, c_0] = Id_{\mathfrak{X}[t_0,c_0]}$. *If* $c_0 \neq -1$, *then*

$$C_4'' := I_{\mathfrak{X}}[t_0; c_0, -c_0]\,(C_4[t_0, c_0])$$

is a new -1*-curve on* $I_{\mathfrak{X}}[t_0, -c_0]$ *with* $(C_4''[t_0, -c_0] \,.\, C_1[t_0, -c_0]) = 2$.

We have to study the curves C_5 and C_6. We have the following lemmas.

Lemma (2.19). *We have* (**i**) $(C_5 \,.\, D_0) = 1$, (**ii**) $(C_5 \,.\, C_1) = 1$, (**iii**) $(C_5 \,.\, C_2) = 0$, (**iv**) $(C_5 \,.\, C_3) = 0$, (**v**) $(C_5 \,.\, C_4) = 3$, (**vi**) $(C_5 \,.\, D_i) = 0$ *for* $2 \leq i \leq 7$, (**vii**) $(C_5 \,.\, D_1) = 1$.

Lemma (2.20). *We have* (**i**) $(C_6 \,.\, D_0) = 1$, (**ii**) $(C_6 \,.\, C_1) = 0$, (**iii**) $(C_6 \,.\, C_2) = 3$, (**iv**) $(C_6 \,.\, C_3) = 1$, (**v**) $(C_6 \,.\, C_4) = 0$, (**vi**) $(C_6 \,.\, D_i) = 0$ *for* $1 \leq i \leq 6$, (**vii**) $(C_6 \,.\, D_7) = 1$.

We prove only Lemma (2.20). Then Lemma (2.19) is proved by a similar method or once we have Lemma (2.20), by the isomorphism $J_{\mathfrak{X}[t_0,c_0]}(c_0, -1-c_0) : \mathfrak{X}[t_0, c_0] \to \mathfrak{X}[t_0, -1-c_0]$ in view of Proposition (1.11) and (2.16).

Proof. The curve $C_6^0 \cap W_4$ on W_4 is defined by

$$2z_4 - y_4^4 + ty_4^2 z_4 + (2c_0+1)y_4^3 z_4 = 0 \tag{2.21.1}$$

by Lemma (2.4). C_6^0 and

$$C_1^0 \cap W_4 = \{(y_4, z_4) \in W_4 \,|\, z_4 = 0\} \tag{2.21.2}$$

have a point $(y_4, z_4) = 0$ in common. Since by (2.19.1) and (2.19.2),

$$z_4 = \frac{y_4^4}{2 + ty_4^2 + (2c_0+1)y_4^3}.$$

The intersection multiplicity of $C_6^0 \cap S$ at $(y_4, z_4) = (0, 0)$ is 4. On the other hand, the curve $C_6^0 \cap W_2$ on W_2 is defined by $2y_2^3 z_2 + y_2 + ty_2 z_2 + (c_0+1)z_2 = 0$ by (2.3.8), which is definition of curve C_6^0. So the intersection multiplicity $C_6^0 \cap S$ at $(y_2, z_2) = (0, 0)$ is 1. Therefore we have

$$\left(C_6^0 \,.\, S\right) = 5. \tag{2.22.1}$$

It follows from (2.19.1) and (2.3.1)

$$\left(C_6^0 \,.\, C_1^0\right) = 1. \tag{2.22.2}$$

Writing $C_6^0 \sim aS + bC_1^0$, we conclude from (2.20.1) and (2.20.2),

$$C_6^0 \sim S + 3C_1^0. \tag{2.23}$$

In particular we have $\left(C_6^0\right)^2 = 8$. Now it follows from Sublemma (3.6) of [U2], $\left(C_6\right)^2 = 0$ because C_6^0 passes through all the centers a_i $(1 \le i \le 8)$. Now $(C_6^0 \,.\, C_2^0) = (S+3C_1^0 \,.\, S-C_1^0) = S^2+2(C_1^0 \,.\, S) = 4$. Since the intersection multiplicity of $C_6^0 \cap C_2^0$ at $(y_4, z_4) = 0$ is 1, we have $(C_6 \,.\, C_2) = 3$. Other multiplicities are calculate in a similar way starting from (2.23).

Theorem (2.24). **(i)** *$L(A_1^{(1)}) := L\left(E_7^{(1)}\right)^{\perp}$ is isomorphic to $\mathbf{Z}^{\oplus 2}$ with the inner product given by minus of Cartan matrix*

$$\begin{bmatrix} -2 & 2 \\ 2 & -2 \end{bmatrix}$$

of the affine root system of type $A_1^{(1)}$.
(ii) *We have $L\left(A_1^{(1)}\right)^{\perp} = L\left(E_7^{(1)}\right)$.*

Proof. We denote by $[C]$ the linear equivalence class of a curve, which we often denote by abuse of notation simply by C. We set

$$M = \langle [C_2] - [C_1], [C_4] - [C_3] \rangle \subset \text{Pic } \mathfrak{X}[t_0, c_0].$$

So M is a subgroup of rank 2 in Pic $\mathfrak{X}[t_0, c_0]$. We have $\left(C_2 - C_1\right)^2 = \left(C_2\right)^2 + \left(C_1\right)^2 = -2$, $\left(C_4 - C_3\right)^2 = \left(C_4\right)^2 + \left(C_3\right)^2 = -2$ by Corollary (2.7), Lemma (2.8), Corollaries (2.12) and (2.14). Moreover $(C_2 - C_1 \,.\, C_4 - C_3) = (C_2 \,.\, C_4) - (C_1 \,.\, C_4) - (C_2 \,.\, C_3) + (C_1 \,.\, C_3) = 2$ by Lemma (2.4). Therefore the inner product on M is given by minus of the Cartan matrix of the affine root system of type $A_1^{(1)}$. It follows from Lemma (2.6), (2.8), Corollaries (2.12), (2.14) $M \subset L\left(E_7^{(1)}\right)^{\perp}$. We have to show $M = L\left(E_7^{(1)}\right)^{\perp}$. Since M and $L\left(E_7^{(1)}\right)^{\perp}$ are of rank 2, we have $\mathbb{Q} \otimes_{\mathbf{Z}} M = \mathbb{Q} \otimes_{\mathbf{Z}} L\left(E_7^{(1)}\right)^{\perp}$. We work in $\mathbb{Q} \otimes_{\mathbf{Z}}$ Pic $\mathfrak{X}[t_0, c_0]$. Let

$$a\,(C_2 - C_1) + b\,(C_4 - C_3) \in \mathbb{Q} \otimes_{\mathbf{Z}} M \subset \mathbb{Q} \otimes_{\mathbf{Z}} \text{Pic } \mathfrak{X}[t_0, c_0] \qquad (a, b \in \mathbb{Q}).$$

To prove (i), we have to show that if

$$a\,(C_2 - C_1) + b\,(C_4 - C_3) \in \text{Pic } \mathfrak{X}[t_0, c_0], \tag{2.25}$$

then $a, b \in \mathbf{Z}$. In fact, by (2.23)

$$(C_2 \,.\, a\,(C_2 - C_1) + b\,(C_4 - C_3)) = -a + 2b \in \mathbf{Z}, \tag{2.26.1}$$

$$(C_4 \,.\, a\,(C_2 - C_1) + b\,(C_4 - C_3)) = 2a - b \in \mathbf{Z}, \tag{2.26.2}$$

$$(C_5 \,.\, a\,(C_2 - C_1) + b\,(C_4 - C_3)) = -a + 3b \in \mathbf{Z}, \tag{2.26.3}$$

$$(C_6 \,.\, a\,(C_2 - C_1) + b\,(C_4 - C_3)) = 3a - b \in \mathbf{Z}. \tag{2.26.4}$$

It follows from (2.26.1) and (2.26.3), $b \in \mathbf{Z}$. Similarly by (2.26.2) and (2.26.4) we conclude $a \in \mathbf{Z}$. Now we show $L\left(A_1^{(1)}\right)^{\perp} = L\left(E_7^{(1)}\right)$. Since $L\left(E_7^{(1)}\right) \subset L\left(A_1^{(1)}\right)^{\perp}$, we may argue as above. Namely $F = \sum_{i=0}^{7} a_i D_i \in \mathbb{Q} \otimes_{\mathbf{Z}} L\left(E_7^{(1)}\right)$ with $a_1 \in \mathbb{Q}$, we have to show that if $F \in \text{Pic } \mathfrak{X}[t_0, c_0]$, then $a_i \in \mathbf{Z}$. Since $(C_1 \,.\, F) \in \mathbf{Z}$ which is equal to a_i by Corollary (2.7) and Lemma (2.8), so $a_1 \in \mathbf{Z}$. Similarly using C_3, we conclude $a_7 \in \mathbf{Z}$. Now $(C_5 \,.\, F) \in \mathbf{Z}$ which is equal to $a_1 + a_0$ by Lemma (2.19) so that $a_1 + a_0 \in \mathbf{Z}$ and consequently $a_0 \in \mathbf{Z}$. Now since

$$(D_1 \,.\, F) = -2a_1 + a_2 \in \mathbf{Z},$$

$$(D_7 \,.\, F) = -2a_7 + a_6 \in \mathbf{Z},$$

we conclude $a_2, a_6 \in \mathbf{Z}$. We have further

$$(D_2 \,.\, F) = a_1 - 2a_2 + a_3 \in \mathbf{Z},$$

$$(D_6 \,.\, F) = a_7 - 2a_6 + a_5 \in \mathbf{Z},$$

so that $a_3, a_5 \in \mathbf{Z}$. Finally

$$(D_3 \,.\, F) = a_2 - 2a_3 + a_4 \in \mathbf{Z}$$

so that $a_4 \in \mathbf{Z}$.

Writing $\mathfrak{X}[t_0, c_0]$ by X, we have an exact sequence of homology

$$\to H_3(X; D, \mathbf{Z}) \to H_2(X \backslash D, \mathbf{Z}) \to H_2(X, \mathbf{Z}) \to H_2(X; D, \mathbf{Z}) \to . \tag{2.27}$$

By the Poincaré duality, we have

$$H_3(X; D, \mathbf{Z}) = H^1(D, \mathbf{Z}) = 0 \tag{2.28}$$

and

$$H_2(X; D, \mathbf{Z}) = H^2(D, \mathbf{Z}) \simeq \bigoplus_{i=0}^{7} \mathbf{Z} D_i. \tag{2.29}$$

Since X is a non-singular, projective rational surface Pic $X \simeq H_2(X, \mathbf{Z})$. So it follows from the exact sequence (2.25) and (2.28), (2.29) that

$$L\left(E_7^{(1)}\right)^{\perp} = H_2(X \backslash D, \mathbf{Z}). \tag{2.30}$$

Let $i : \mathbb{C}(t, c,) \to \mathbb{C}(t, c)$ be the $\mathbb{C}(t)$-automorphism of the field $\mathbb{C}(t, c)$ sending c to $-c$. Similarly, let $j : \mathbb{C}(t, c) \to \mathbb{C}(t, c)$ be the $\mathbb{C}(t)$-automorphism of the field $\mathbb{C}(t, c)$ such that $j(c) = -1 - c$. So the subgroup $< i, j >$ of the automorphisms group of the field $\mathbb{C}(t, c)$ generated by i and j is isomorphic to the extended affine Weyl group $\widehat{G}$. The group W_a operates on Spec $\mathbb{C}(t, c)$. The group $\widehat{G}$ also operates on $\mathfrak{X}_{\mathbb{C}(t,c)} := \mathfrak{X} \otimes_{\mathbb{C}[t,c]} \mathbb{C}(t, c)$ in such a way that the morphism $\mathfrak{X}_{\mathbb{C}(t,c)} \to$ Spec $\mathbb{C}(t, c)$ is $\widehat{G}$-equivariant. A -1-curve C on a surface is a curve isomorphic to $\mathbb{P}^1$ with $C^2 = -1$.

Theorem (2.31). *For every $(t_0, c_0) \in \mathbb{C}^2$, there are infinitely many -1-curves on $\mathfrak{X}[t_0, c_0]$.*

Proof. First we assume that c_0 is not an integer. Then we have an isomorphism

$$T_+[t_0\,;\, c_0 - 1\,,\, c_0] = I_{\mathfrak{X}}[t_0\,;\, -c_0\,,\, c_0] \circ J_{\mathfrak{X}}[t_0\,;\, c_0 - 1\,,\, -c_0] : \mathfrak{X}[t_0\,,\, c_0 - 1] \to \mathfrak{X}[t_0\,,\, c_0].$$

So in particular the isomorphism

$$T_+[t_0\,;\, c_0 - 1\,,\, c_0] : X[t_0\,,\, c_0 - 1] \to \mathfrak{X}[t_0\,,\, c_0]$$

maps the -1-curve $C_3[t_0\,,\, c_0 - 1]$ on $\mathfrak{X}[t_0\,,\, c_0 - 1]$ to a -1-curve

$$\Gamma_1[t_0\,,\, c_0] := T_+[t_0\,;\, c_0 - 1\,,\, c_0](C_3[t_0\,,\, c_0 - 1])$$

on $\mathfrak{X}[t_0\,,\, c_0]$. For a positive integer n, the iteration

$$T_+[t_1\,;\, c_0 - n\,,\, c_0] := T_+[t_0\,;\, c_0 - 1\,,\, c_0] \circ T_+[t_0\,;\, c_0 - 2\,,\, c_0 - 1]$$

$$\circ \cdots \circ T_+[t_0\,;\, c_0 - n + 1\,,\, c_0 - n + 2] \circ T_+[t_0\,;\, c_0 - n\,,\, c_0 - n + 1]$$

$$: \mathfrak{X}[t_0\,,\, c_0 - n] \to \mathfrak{X}[t_0\,,\, c_0 - n + 1] \to \cdots \to \mathfrak{X}[t_0\,,\, c_0 - 1] \to \mathfrak{X}[t_0\,,\, c_0]$$

of isomorphisms maps the -1-curve $C_3[t_0\,,\, c_0 - n]$ to a -1-curve $\Gamma_n[t_0\,,\, c_0] := T_+[t_0\,;\, c_0 - n, c_0](C_3[t_0\,,\, c_0 - n])$. Hence we have -1-curves $\Gamma_n[t_0\,,\, c_0]$ $(n = 1, 2, \ldots)$, on $\mathfrak{X}[t_0\,,\, c_0]$. We have to show that the -1-curves $\Gamma_n[t_0\,,\, c_0]$ are distinct curves on $\mathfrak{X}[t_0\,,\, c_0]$. To this end, it is sufficient to show that the linear equivalence classes $[\Gamma_n] \in$ Pic $\mathfrak{X}[t_0\,,\, c_0]$ are distinct. It follows from the construction of $\mathfrak{X}[t_0, c_0]$ that

$$\text{Pic } \mathfrak{X}[t_0\,,\, c_0] = \bigoplus_{i=0}^{7} \mathbb{Z}D_i \bigoplus \mathbb{Z}C_1 \bigoplus \mathbb{Z}C_3.$$

By Proposition (2.16) and (2.17), we have

$$T_+[t_0\,;\,c_0-1\,,\,c_0]\,(C_1[t_0\,,\,c_0-1]) = C_3[t_0\,,\,c_0],$$

$$T_+[t_0\,;\,c_0-1\,,\,c_0]\,(C_3[t_0\,,\,c_0-1]) = C_2[t_0\,,\,c_0]. \tag{2.32}$$

Let us express the linear equivalence class of C_2 as a linear combination of C_1, C_3 and the D_i's. Namely we set

$$C_2 \sim a_1C_1 + a_3C_3 + \sum_{i=0}^{7} b_iD_i$$

and determine the integers a_1, a_3 and b_j's. We have linear equations

$$0 = (C_1\,,\,C_2) = -a_1 + b_1,$$

$$1 = (D_1\,,\,C_2) = a_1 - 2b_1 + b_2,$$

$$0 = (D_2\,,\,C_2) = b_1 - 2b_2 + b_3,$$

$$0 = (D_3\,,\,C_2) = b_2 - 2b_3 + b_4,$$

$$0 = (D_4\,,\,C_2) = b_3 - 2b_4 + b_5 + b_0,$$

$$0 = (D_0\,,\,C_2) = -2b_0 + b_4,$$

$$0 = (D_5\,,\,C_2) = b_4 - 2b_5 + b_6,$$

$$0 = (D_6\,,\,C_2) = b_5 - 2b_6 + b_7,$$

$$0 = (D_7\,,\,C_2) = -2b_7 + b_6 + a_3,$$

$$0 = (C_3\,,\,C_2) = -b_7 - a_3.$$

We solve this system of linear equations to get

$$\Gamma_1 = C_2 \sim -C_1 + 2C_3 - D_1 + D_3 + 2D_4 + 2D_5 + 2D_6 + 2D_7 + D_0.$$

In particular, we have

$$\Gamma_1 = T_+[t_0\,;\,c_0-1\,,\,c_0](C_3) \equiv -C_1 + 2C_3 \tag{2.33}$$

$$T_+[t_0\,;\,c_0-1\,,\,c_0](C_1) \equiv C_3 \tag{2.34}$$

modulo the subgroup $L\left(E_7^{(1)}\right) = \langle D_0\,,\, D_1\,,\, \ldots\,,\, D_7\rangle$ in Pic $X[t_0\,,\, c_0]$. Since $L\left(E_7^{(1)}\right)$ is invariant under $I_{\mathfrak{X}}$ and $J_{\mathfrak{X}}$ by Proposition (1.10) and hence by T_+, it follows from (2.34) and (2.35),

$$\begin{aligned}
\Gamma_2 &= T_+[t_0\,;\, c_0-1\,,\, c_0]\circ T_+[t_0\,;\, c_0-2\,,\, c_0-1]\,(C_3[t_0\,,\, c_0-2]) \\
&\equiv T_+[t_0\,;\, c_0-1\,,\, c_0]\,(-C_1[t_0\,,\, c_0-1] + 2C_3[t_0\,;\, c_0-1]) \\
&\equiv -C_3[t_0\,,\, c_0] + 2\,(-C_1[t_0\,,\, c_0] + 2C_3[t_0\,;\, c_0]) \\
&\equiv -2C_1[t_0\,,\, c_0] + 3C_3[t_0\,;\, c_0]
\end{aligned}$$

modulo $\Gamma\left(E_7^{(1)}\right)$. So Γ_1 and Γ_2 are distinct curves on $X[t_0\,,\, c_0]$. Similarly we have

$$\begin{aligned}
\Gamma_3 &\equiv -6C_1 + 10C_3 \\
\Gamma_4 &\equiv -20C_1 + 34C_3 \\
\Gamma_5 &\equiv -34C_1 + 116C_3 \\
&\vdots
\end{aligned}$$

so that Γ_1, Γ_2, $\cdots$ are distinct -1-curves on $\mathfrak{X}[t_0\,,\, c_0]$. If c_0 is an integer, since the surface $\mathfrak{X}[t_0\,,\, c_0]$ is isomorphic to $\mathfrak{X}[t_0, -1]$ as we proved in [U2], we may assume $c_0 = -1$. Then we can apply the above argument.

§3. Vanishing cycles on $\mathfrak{X}[t_0\,,\, c_0]$.

If $c_0 = 1$, then we have the -2-curve $C_2'[t_0\,,\, 0]$ on $\mathfrak{X}[t_0, 0]$ that is the limit of the cycle $C_2[t_0, c] - C_1[t_0, c]$ on $\mathfrak{X}[t_0, c]$ when $c \to 0$. We can collapse the -2-curve $C_2'[t_0, 0]$ to a normal singular point to get a normal algebraic surface $\mathfrak{X}''[t_0]$ with an ordinary double point. This suggests that the cycle $C_2[t_0, c] - C_1[t_0, c]$ on $\mathfrak{X}[t_0, c]$ is a vanishing cycle.

Proposition (3.1). *There exists a flat family* $\varphi' : \mathfrak{X}'[t, c^2] \to \operatorname{Spec} \mathbb{C}[t, c^2]$ *of projective algebraic surfaces with the following properties.*
(i) *The variety* $\mathfrak{X}'[t, c^2]$ *is non-singular.*
(ii) *We have an isomorphisms*

$$\mathfrak{X}[t_0, c_0] \simeq \mathfrak{X}'[t_0, c_0^2] \qquad \textit{if } c_0 \neq 0,$$

and

$$\mathfrak{X}''[t_0] \sim \mathfrak{X}'[t_0, 0].$$

Here $\mathfrak{X}'[t_0, c_0^2]$ *denotes the fiber* ${\varphi'}^{-1}((t_0, c_0^2))$ *over a point* $(t_0, c_0^2) \in \mathrm{Spec}\ \mathbb{C}[t, c^2]$.
(iii) *The morphism* $\varphi : \mathfrak{X}'[t, c^2] \to \mathrm{Spec}\ \mathbb{C}[t, c^2]$ *is smooth except for the points on* $\mathfrak{X}'[t_0, 0]$ *that correspond to the rational double point of* $\mathfrak{X}''[t_0]$*,* $t_0 \in \mathbb{C}$*.*
(iv) *We have a* $\mathbb{C}[t, c, 1/c]$*-isomorphism*

$$\mathfrak{X}[t, c] \otimes_{\mathbb{C}[t,c]} \mathbb{C}\left[t, c, \frac{1}{c}\right] \simeq \mathfrak{X}'[t, c^2] \otimes_{\mathbb{C}[t,c^2]} \mathbb{C}\left[t, c, \frac{1}{c}\right].$$

Proof. We consider a $\mathbb{C}[t, c]$-morphism

$$\begin{aligned} f_1 \quad &: \quad W_1 \times \mathrm{Spec}\ \mathbb{C}[t, c] = W_1 \times \mathbb{C}^2 \to \mathbb{P}^3 \times \mathbb{C}^2 \\ & \qquad (y_1, t_1; t, c) \mapsto (1, y_1(c - y_1 z_1), 2y_1 z_1 - c, z_1; t, c). \end{aligned}$$

The morphism f_1 extends to a rational map

$$f : Z_0[t, c] \to \mathbb{P}^3 \times \mathbb{C}^2,$$

which turns out as we can check it easily, to be a morphism. We recall that $Z_0[t, c]$ is the starting family of ruled surfaces in the construction of $\mathfrak{X}[t, c]$. For example, we have on $W_3 \times \mathrm{Spec}\ \mathbb{C}[t, c]$,

$$\begin{aligned} f_3 \quad &: \quad W_3 \times \mathbb{C}^2 \to \mathbb{P}^3 \times \mathbb{C}^2 \\ & \qquad (y_3, z_3; t, c) \mapsto (1, z_3, c - 2y_3 z_3, y_3(c - y_3 z_3); t, c). \end{aligned}$$

Moreover f_3 factors through a family of quadratic surface

$$Q[t, c^2] : 4x_1 x_3 - x_2^2 + c^2 x_0^2 = 0$$

in $\mathbb{P}^3 \times \mathbb{C}^2$ so that the surface $Q[t, c^2]$ is defined over $\mathbb{C}[t, c^2]$. If $c = 0$, f_3 collapses the curve $C_2'[t_0, 0]$ for every t_0. We can check that the morphism $f : Z_0[t, c] \to Q[t, c^2]$ is an isomorphism outside the curve $C_2'[t, 0]$. We can also show by calculation that we can blow up $Z_0[t, c]$ and $Q[t, c^2]$ so that we have a morphism

$$\widetilde{f} : \mathfrak{X}[t, c] \to \widetilde{Q}[t, c^2] =: \mathfrak{X}'[t, c^2]$$

(cf. Proposition (1.11)). Then $\widetilde{f}$ has the required properties. We proved in [U2] that we have an isomorphism

$$I_{\mathfrak{X}}[t; c, -c] : \mathfrak{X}[t, c] \otimes_{\mathbb{C}[t,c]} \mathbb{C}[t, c, 1/c] \to \mathfrak{X}[t, c] \otimes_{\mathbb{C}[t,c]} \mathbb{C}[t, c, 1/c]$$

covering Spec $\mathbb{C}[t,c] = \mathbb{C}^2 \to$ Spec $[t,c] = \mathbb{C}^2$, $(t,c) \mapsto (t,-c)$. The quotient variety

$$\begin{array}{c} \mathfrak{X}[t,c] \otimes_{\mathbb{C}[t,c]} \mathbb{C}[t,c,1/c] / \left\langle I_{\mathfrak{X}}[t;c,-c] \right\rangle \\ \downarrow \\ \text{Spec } \mathbb{C}[t,c,1/c]^{<I>} = \text{Spec } \mathbb{C}[t,c^2,1/c^2] \end{array}$$

is isomorphic to

$$\begin{array}{c} \mathfrak{X}'[t,c^2] \otimes_{\mathbb{C}[t,c^2]} \mathbb{C}[t,c^2,1/c^2] \\ \downarrow \\ \text{Spec } \mathbb{C}[t,c^2,1/c^2]. \end{array}$$

This shows that the involution $I_{\mathfrak{X}}[t;c,-c]$ arises form the rational double point of type A_1. Namely if we fix $t = t_0 \in \mathbb{C}$, the birational involution

$$I : \mathfrak{X}[t_0,c] \to \mathfrak{X}[t_0,c] \tag{3.2}$$

comes from the automorphism of the singular quadratic surface

$$Q = \left\{ (x_0,x_1,x_2;c) \in \mathbb{P}^2 \times \mathbb{C} \,|\, 4x_1x_3 - x_2^2 + c^2x_0 = 0 \right\}$$

sending $(x_0,x_1,x_2;c)$ to $(x_0,x_1,x_2;-c)$. So the birational map is the simplest example of flop between 3 dimensional varieties and hence this is why I is not an isomorphism (cf. [U2] and [K], 12.1. Example). Similarly the cycle $C_4 - C_3$ is the vanishing cycle around $c = -1$.

§4. Regular functions on $\mathfrak{X}[t_0,c_0] \backslash D$.

We begin with a lemma.

Lemma (4.1). *If there exists a complete curve C on $\mathfrak{X}[t_0,c_0] \backslash \mathfrak{D}[t_0,c_0]$, then $C \cap W_1 \subset W_1$ is an open algebraic curve.*

Proof. It follows from the construction of the surface $\mathfrak{X}[t_0,c_0]$ that $\mathfrak{X}[t_0,c_0] \backslash W_1 = \mathfrak{D}[t_0,c_0] \cup C_1 \cup C_4$. The curve C_1 intersects $\mathfrak{D}[t_0,c_0]$ on D_1 and the curve C_4 intersects $\mathfrak{D}[t_0,c_0]$ on D_7.

Proposition (4.2). (i) *If the parameter c_0 is not an integer, then there is no projective algebraic curve on $\mathfrak{X}[t_0,c_0] \backslash \mathfrak{D}[t_0,c_0]$.* **(ii)** *If c_0 is an integer, then there is only one projective algebraic curve C on $\mathfrak{X}[t_0,c_0] \backslash \mathfrak{D}[t_0,c_0]$. The curve C is isomorphic to $\mathbb{P}^1$ and $C^2 = -2$. In other words, C is a -2-curve.*

Proof. Let us assume that there exists a complete curve C on $\mathfrak{X}[t_0,c_0] \backslash \mathfrak{D}[t_0,c_0]$. Since the Painlevé equations have no movable singular points and since $\mathfrak{X}[t_0,c_0] \backslash \mathfrak{D}[t_0,c_0]$

is the space of initial conditions, integrating the curve C along the foliation on $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ generated by the Hamiltonian system

$$S_2(c_0) \begin{cases} \dfrac{dy_1}{dt} = y_1^2 + z_1 + \dfrac{t}{2}, \\ \\ \dfrac{dz_1}{dt} = -2y_1 z_1 + c_0 \end{cases}$$

on W_1, we get an analytic family $\mathfrak{C} \to (\text{Spec } \mathbb{C}[t])^{an}$ of complete algebraic curves on $\mathfrak{X}[t, c_0] \to \text{Spec } \mathbb{C}[t]$. So by G.A.G.A., there exists a field extension $K \supset \mathbb{C}(t)$ such that K is a subfield of the field of meromorphic functions on an open set of $\mathbb{C}$ with coordinate system t and such that the analytic family $\mathfrak{C} \to (\text{Spec } \mathbb{C}[t])^{an}$ is an algebraic family of curves defined over the field K. We may assume that the field is closed under the derivation d/dt. Let $F(y_1, z_1) = 0$ be a defining equation of $(W_1 \times \mathbb{C} \cap \mathfrak{C}) \to \text{Spec } K$ on $(\mathfrak{X}[t, c_0] \cap W_1) \otimes_{\mathbb{C}[t]} K \to \text{Spec } K$ so that $F(y_1, z_1) \in K[y_1, z_1]$. Since $W_1 \times \mathbb{C} \cap \mathfrak{C}$ is invariant under the Hamiltonian flow

$$D(c_0) = \frac{\partial}{\partial t} + \left(y_1^2 + z_1 + \frac{t}{2}\right) \frac{\partial}{\partial y_1} + (c_0 - 2y_1 z_1) \frac{\partial}{\partial z_1},$$

$f(y_1, z_1)$ divides $D(c)\,(f(y_1, z_1))$ in the polynomial ring $K[y_1, z_1]$. In the language of [U1], F is an invariant polynomial over K. Then Theorem (2.1), [UW] implies $c_0 = \alpha_0 - \frac{1}{2} \in \mathbf{Z}$ and the curve C arises from the Riccati equation and $C^2 = -2$.

Lemma (4.3). *The canonical divisor K of $\mathfrak{X}[t_0, c_0]$ is given by a divisor $-F$, where*

$$F = 2D_0 + D_1 + 2D_2 + 3D_3 + 4D_4 + 3D_5 + 2D_6 + D_7. \tag{4.3.1}$$

Proof. The lemma follows from the following two observations and the construction of $\mathfrak{X}[t_0, c_0]$. First, the canonical divisor K_0 of $Z_0[t_0, c_0]$ is given by $-2S$. Second, let K_i be canonical divisor of $Z_i[t_0, c_0]$ $(0 \le i \le 8)$. Then $K_{i+1} \sim \pi_{i+1}^* K_i + E_{i+1}$, where E_i is exceptional divisor of $\pi_{i+1} : Z_{i+1}[t_0, c_0] \to Z_i[t_0, c_0]$ $0 \le i \le 7$.

F is the null vector of the Cartan matrix C of $\widetilde{A}_1$. Namely $C^t(2, 1, 2, 3, 4, 3, 2, 1) = 0$.

Proposition (4.4). $h^0\left(\mathfrak{X}[t_0, c_0], -K\right) = 1$.

Proof. Since F is effective, $h^0\left(\mathfrak{X}[t_0, c_0], -K\right) \ge 1$. Assume $h^0\left(\mathfrak{X}[t_0, c_0], -K\right) \ge 2$. Then there exists an effective divisor H linearly equivalent to $-K$ and distinct of F. Let $H = H_1 + H_2$ be a decomposition into two effective divisors such that the support of H_1 is a subset of $\{D_i \,|\, 0 \le i \le 7\}$ and such that no irreducible component of H_2 is D_i, $0 \le i \le 7$. We show $H_2 = 0$. Let us assume $H_2 \ne 0$. Since we blow up the ruled surface $Z_0[t_0, c_0]$ 8 times,

$$0 = K^2 = (K\,.\,-H) = (K\,.\,-H_1 - H_2) = -(K\,.\,H_1) - (K\,.\,H_2). \tag{4.5}$$

Since D_i is a -2-curve, $(K\,.\,D_i) = 0$ for $0 \leq i \leq 7$. So

$$(4.6) \qquad (K\,.\,H_1) = 0.$$

First assume that c_0 is not an integer. Then there is no projective curve on $\mathfrak{X}[t_0, c_0] \backslash \bigcup_{i=0}^{7} D_i$ by Lemma (4.1). Hence $(-K\,.\,H_2) = (F\,.\,H_2) > 0$. This contradicts (4.5) and (4.6). Therefore $H_2 = 0$ or the support of $H \subset \bigcup_{i=0}^{7} D_i$. Since the D_i's are linearly independent divisors in Pic $\mathfrak{X}[t_0, c_0]$, this shows $H = F$, which is contradiction. If c_0 is an integer, there exists the unique projective curve C on $\mathfrak{X}[t_0, c_0] \backslash \bigcup_{i=0}^{7} D_i$. If we assume $H_2 \neq 0$, then the above argument shows $H_2 = nC$ for an appropriate integer $n \geq 0$. Since C is a -2-curve,

$$0 = (K.C) = (-H.C) = (-H_1 - H_2.C) = (-H_1.C) - (H_2.C) = -n(C.C) = 2n.$$

So we conclude $H_1 = 0$ and the argument above leads us to a contradiction.

Corollary (4.7). (to the proof.) *For every integer $m \geq 0$,*

$$H^0(\mathfrak{X}[t_0, c_0], -mK) = \mathbb{C}.$$

Proof. The assertion being trivial for $m = 0$, we may assume $m > 0$. Then the argument of the proof of Proposition (4.4) allows us to prove the corollary. We can formulate Corollary (4.7) in another form.

Corollary (4.8).

$$H^0(\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0], \mathcal{O}) = \mathbb{C}.$$

Proof. Since $-K \sim 2D_0 + D_1 + 2D_2 + 3D_3 + 4D_4 + 3D_5 + 2D_6 + D_7$, this result follows from Corollary (4.7).

§5. Takano coordinate systems.

It follows from Proposition (4.4) that there exists a 2-form ω on $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ such that ω vanishes at no point of $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$. Moreover ω is unique up to a non-zero constant multiplication. Namely, there exists a symplectic structure on $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$. K. Takano and his colleagues introduced nice coordinate systems on $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ such that locally the 2-form is written in a canonical form in terms of the coordinate system. They cover $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ by there copies of $\mathbb{C}^2$ such that coordinate transformations are symplectic. We call these coordinate systems the Takano coordinate systems. To explain the Takano coordinate systems, let us come back to the construction of $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$. To construct $\mathfrak{X}[t_0, c_0]$, we started from the ruled surface $Z_0[t_0, c_0]$. We blow up $Z_0[t_0, c_0]$ eight times and removed the proper transform D_0 of S, the proper transforms $D_1, D_2, \ldots, D_7$ of all the exceptional divisor except for the last exceptional divisor $E_8 = C_3 = D_8$. Let us recall

$$S = \{(y_2, z_2) \in W_2 \,|\, z_2 = 0\} \cup \{(y_4, z_4) \in W_4 \,|\, z_4 = 0\}$$

so that

$$Z_0[t_0, c_0] = W_1 \cup W_3 \cup S.$$

Hence

$$\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0] = W_1 \cup W_3 \cup (D_8 \backslash (D_8 \cap D_7)) \tag{5.1}$$

(cf. Fig.(1.3)). So we start from $W_1 \cup W_3 \cup W_4$. We use the notation of §2, [U2]. We have to resolve the rational map

$$F : W_4 \cdots \to \mathbb{P}^1, \quad (y_4, z_4) \mapsto (y_4^4 z_4, 2z_4 - y_4^4 + t y_4^2 z_4 + (2c+1) y_4^3 z_4) \tag{5.2}$$

(cf. Sublemma (3.6) in [U2]). Let (y_{4+i}, z_{4+i}) be the coordinate system of $W_{4+i}(z) = \mathbb{C}^2$ $1 \leq i \leq 4$ so that we have $y_{4+i} = y_4$, $z_{4+i-1} = y_{4+i} z_{4+i}$ $(1 \leq i \leq 4)$. Hence

$$y_4 = y_5 = y_6 = y_7 = y_8, \quad y_4 = y_8^4 z_8. \tag{5.3}$$

On $W_8(z)$, F is written as

$$F : W_8(z) \to \mathbb{P}^1, \quad (y_8^4 z_8, 2z_8 - 1 + t y_8^2 z_8 + (2c+1) y_8^3). \tag{5.4}$$

Now we introduce a coordinate system (y_8, v_8) on $W_8(z)$ such that

$$v_8 = \frac{1}{z_8}. \tag{5.5}$$

Precisely speaking, we consider $W_8'(z) = \mathbb{A}^2$ with coordinate system (y_8, z_8) and birational map

$$W_8'(z) \cdots \to W_8(z), \quad (y_8, v_8) \mapsto (y_8, 1/v_8). \tag{5.6}$$

Let $W_8'(z)^0 = \{(y_8, v_8) \in W_9'(z) \,|\, v_8 \neq 0\}$ so that $W_8'(z)^0$ is identified with an open set of $W_8(z)$ by (5.6). We have on $W_8'(z)$

$$F : W_8'(z) \to \mathbb{P}^1, \quad (y_8, v_8) \mapsto (y_8^4, 2 - v_8 + t y_8^2 + (2c+1) y_8^3). \tag{5.7}$$

So $(y_8, v_8) = (0, 2)$ is the singular point of F in (5.7). We blow up $W_8'(z)$ at $(y_8, v_8) = (0, 2)$ to resolve the rational map F given by (5.7). We set

$$v_8 - 2 = y_9 z_9, \quad y_8 = y_9. \tag{5.8}$$

On $W_9 = \mathbb{A}^2$ with coordinate system (y_9, z_9), we have

$$F : W_9 \to \mathbb{P}^1, \quad (y_9, z_9) \mapsto (y_9^3, -z_9 + ty_9 + (2c+1)y_9^2). \tag{5.9}$$

The singular point of F in (5.9) is $(y_9, z_9) = (0, 0)$. We blow up W_9 at this point. So we set

$$y_{10} = y_9, \quad z_9 = y_9 z_{10}. \tag{5.10}$$

On $W_{10} = \mathbb{A}^2$ with coordinate system (y_{10}, z_{10}), we have

$$F : W_{10} \to \mathbb{P}^1, \quad (y_{10}, z_{10}) \mapsto (y_{10}^2, -z_{10} + t + (2c+1)y_{10}). \tag{5.11}$$

The singular point of F in (5.11) is $(y_{10}, z_{10}) = (0, t)$. We blow up W_{10} at (0,t). So we set

$$y_{11} = y_{10}, \quad z_{10} - t = y_{11} z_{11}. \tag{5.12}$$

On $W_{11} = \mathbb{A}^2$ the coordinate system (y_{11}, z_{11}), we have

$$W_{11} \to \mathbb{P}^1, \quad (y_{11}, z_{11}) \mapsto (y_{11}, -z_{11} + (2c+1)). \tag{5.13}$$

The singular point of (5.11) is $(y_{11}, z_{11}) = (0, 2c+1)$. We blow up W_{11} at $(0, 2c+1)$. So we set

$$y_{12} = y_{11}, \quad z_{11} - (2c+1) = y_{12} z_{12}. \tag{5.14}$$

On $W_{12} = \mathbb{C}^2$ with coordinate system (y_{12}, z_{12}), we have

$$F : W_{12} \to \mathbb{P}^1, \quad (y_{12}, z_{12}) \mapsto (1, z_{12}). \tag{5.15}$$

Namely the rational map is resolved on W_{12}. It follows from (5.8), (5.10), (5.12), (5.14)

$$v_8 = y_{12}^4 z_{12} + (2c+1)y_{12}^3 + ty_{12}^2 + 2. \tag{5.16}$$

So we have a rational map

$$W_{12} \cdots \to W_8'(z) \to W_8(z) \tag{5.17}$$

sending (y_{12}, z_{12}) to

$$(y_8, z_8) = \left(y_{12}, \frac{1}{y_{12}^4 z_{12} + (2c+1)y_{12}^3 + ty_{12}^2 + 2} \right).$$

We have to restrict the map (5.17) on an open set

$$W_{12}^0 = \left\{ (y_{12}, z_{12}) \,|\, y_{12}^4 z_{12} + (2c+1)y_{12}^3 + ty_{12}^2 + 2 \neq 0 \right\}.$$

Then we have a regular map

$$W_{12}^0 \to W_8(z), \tag{5.18}$$

$$(y_{12}, z_{12}) \mapsto (y_8, z_8) = \left(y_{12}, \frac{1}{y_{12}^4 z_{12} + (2c+1)y_{12}^3 + ty_{12}^2 + 2} \right).$$

Hence it follows from (5.3), (5,18) that we have a regular map

$$W_{12}^0 \to W_4(z), \tag{5.19}$$

$$(y_{12}, z_{12}) \mapsto (y_8, z_8) = \left(y_{12}, \frac{y_{12}^4}{y_{12}^4 z_{12} + (2c+1)y_{12}^3 + ty_{12}^2 + 2} \right).$$

Since $z_4 = \frac{1}{z_3}$, we have by (5.19)

$$\frac{1}{z_3} = \frac{y_{12}^4}{y_{12}^4 z_{12} + (2c+1)y_{12}^3 + ty_{12}^2 + 2}$$

or

$$z_3 = z_{12} + \frac{2c+1}{y_{12}} + \frac{t}{y_{12}^2} + \frac{2}{y_{12}^4}. \tag{5.20}$$

So $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ is covered by W_1, W_3 and W_{12}^0. A point $(y_3, z_3) \in W_3$ and a point $(y_{12}, z_{12}) \in W_{12}^0$ are identified if **(i)** $y_4 = y_{12} \neq 0$ and if **(ii)** we have (5.20). In view of (5.18), we may enlarge W_{12}^0 or we may replace it by W_{12}. So in view of (5.1), we have proved the following

Theorem (5.21). *$\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$ is covered by three copies W_1, W_3, W_{12} of $\mathbb{A}^2$. We glue together these copies by following rule.*
(i) *A point $(y_1, z_1) \in W_1$ and a point (y_3, z_3) are identified if $y_1 y_3 = 1$ and if $z_1 = y_3(c - y_3 z_3)$.*
(ii) *A point $(y_1, z_1) \in W_1$ and a point $(y_{12}, z_{12}) \in W_{12}$ are identified if $y_1 y_{12} = 1$ and if $z_1 + 2y_1^2 + t_0 = y_{12}\left(-(c+1) - y_{12} z_{12}\right)$.*
(iii) *A point $(y_3, z_3) \in W_3$ and a point $(y_{12}, z_{12}) \in W_{12}$ are identified if $y_3 = y_{12} \neq 0$ and if we have*

$$z_3 = z_{12} + \frac{2c+1}{y_{12}} + \frac{t}{y_{12}^2} + \frac{2}{y_{12}^4}.$$

Proof. It is sufficient to see that (ii) is a consequence of (i) and (iii). We can check this by an easy calculation.

Corollary (5.22). *2-forms $dy_1 \wedge dz_1$ on W_1, $dy_3 \wedge dz_3$ on W_3 and $dy_{12} \wedge dz_{12}$ on W_{12} coincide on the intersections $W_i \cap W_j$ and thus define a symplectic structure on $\mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$.*

Proof. This is an immediate consequence of Theorem (5.21).

Remark (5.23). *If we notice the relation (iii) follows from (i) and (ii), we have a symmetry*

$$y_1 \mapsto -y_1, \quad z_1 \mapsto -(z_1 + 2y_1^2 + t), \quad c \mapsto -(c+1),$$

which sends as a consequence $z_1+2y_1^2+t \mapsto -z_1$, $-(c+1) \mapsto -c$, *of* $W_1 \cup W_3 \cup W_{12} = \mathfrak{X}[t_0, c_0] \backslash \mathfrak{D}[t_0, c_0]$.

The open surface $\mathfrak{X}[t_0, c_0]\backslash D$ is covered by three copies W_1, W_3, W_{12} of $\mathbb{A}^2$. As Corollary (5.23) shows, the 2-forms $dy_1 \wedge dz_1$, $dy_3 \wedge dz_3$, $dy_{12} \wedge dz_{12}$ glue together and define a 2-form ω on $\mathfrak{X}[t_0, c_0]\backslash D$. The vanishing cycles span $L\left(A_1^{(1)}\right) = L\left(E_7^{(1)}\right)^{\perp}$ and hence by (2.30) $\mathbf{Z}(C_2 - C_1) \oplus \mathbf{Z}(C_4 - C_3) = H_2(\mathfrak{X}[t_0, c_0]\backslash D, \mathbf{Z})$.

Proposition (5.24).

$$\int_{C_2 - C_1} \omega = c, \quad \int_{C_4 - C_3} \omega = -c - 1.$$

Proof. We show the first equality. If $c = 0$, $C_2 - C_1$ is a complete curve on $\mathfrak{X}[t_0, c_0]\backslash D$ so that

$$\int_{C_2 - C_1} \omega = 0$$

because ω is a holomorphic 2-form. So we may assume $c \neq 0$. Since $y_3 = y_4$, $z_3 = 1/z_4$, we have on W_4

$$\omega = dy_3 \wedge dz_3 = -dy_4 \wedge \frac{1}{z_4^2} dz_4.$$

Blowing up W_4 at $(y_4, z_4) = (0, 0)$, we get a morphism $\widetilde{W}_4 \to W_4$. The surface $\widetilde{W}_4$ is covered by $W_4(y)$ and $W_4(z)$ that are isomorphic to $\mathbb{A}^2$. We have coordinate systems (Y, z) on $W_4(y) \simeq \mathbb{A}^2$ and (y, Z) on $W_4(z) \simeq \mathbb{A}^2$. We identify a point $(Y, z) \in W_4(y)$ and $(y, Z) \in W_4(z)$ if $y = Yz$, $z = yZ$ and if $YZ = 1$. The morphism $\widetilde{W}_4 = W_4(y) \cap W_4(z) \to W_4$ is defined respectively on $W_4(y)$ by

$$W_4(y) \to W_4 = \mathbb{A}^2, \qquad (Y, z) \mapsto (Yz, z)$$

and on $W_4(z)$ by

$$W_4(y) \to W_4 = \mathbb{A}^2, \qquad (y, Z) \mapsto (y, yZ).$$

The curves C_1 and C_2 are defined on W_4 respectively by

$$C_1 \cap W_4 = \left\{(y_4, z_4) \in \mathbb{A}^2 \,|\, y_4 = 0\right\}$$

and

$$C_2 \cap W_4 = \left\{(y_4, z_4) \in \mathbb{A}^2 \,|\, y_4 - cz_4 = 0\right\}.$$

So we have on $W_4(y) \simeq \mathbb{A}^2$,

$$C_1 \cap W_4(y) = \{(Y, z) \in W_4(y) \,|\, Y = 0\}$$

and

$$C_2 \cap W_4(y) = \{(Y, z) \in W_4(y) \,|\, Y = c\}.$$

Now $D_1 \cap W_1(y) = \{(Y, z) \in \mathbb{A}^2 \,|\, z = 0\}$ and hence on $W_4(y)$, $C_1 \cap D_1 = \{(0, 0)\}$, $C_2 \cap D_1 = \{(c, 0)\}$. Now let γ be a segment in $D_1 \cap W_4(y) = \{(Y, z) \in W_4(y) \,|\, z = 0\}$ joining the points $(c, 0)$ and $(0, 0)$. Let τ be a closed tublar neighborhood of γ. We set $C_1 \cap \tau = \tau_1$, $C_2 \cap \tau = \tau_2$. We have on $W_4(y)$

$$-dy_4 \wedge \frac{1}{z_4^2} dz_4 = -dY \wedge \frac{dz}{z}.$$

Since $\partial\tau$ is homologous to $C_2 - C_1$ in $\mathfrak{X}[t_0, c_0] \backslash D$,

$$\int_{C_2 - C_1} \omega = \int_{\partial\tau} \omega = \int_{\partial\tau} -dY \wedge \frac{dz}{z} = \int_{\gamma} -2\pi i dY = 2\pi i c.$$

The isomorphism

$$J(c_0, -c_0 - 1, t_0) : \mathfrak{X}[t_0, c_0] \to \mathfrak{X}[t_0, -c_0 - 1]$$

maps $C_3[t_0, c_0]$ to $C_1[t_0, -c_0 - 1]$, $C_4[t_0, c_0]$ to $C_2[t_0, -c_0 - 1]$ and $\omega[t_0, c_0]$ to $\omega[t_0, -c_0 - 1]$. So

$$\int_{C_4[t_0,c_0] - C_3[t_0,c_0]} \omega[t_0, c_0] = \int_{C_2[t_0,-c_0-1] - C_1[t_0,-c_0-1]} \omega[t_0, -c_0 - 1] = -c_0 - 1.$$

In particular, since either $c \neq 0$ or $-c - 1 \neq 0$, we have proved the following

Corollary (5.25). *The de Rham class of ω in $H^2(\mathfrak{X}[t_0, c_0] \backslash D, \mathbb{C})$ is not 0. In other words the closed 2-form ω on $x[t_0, c_0] \backslash D$ is not exact.*

Theorem (5.26). (Cf. [A]). *We have*

$$H^0(\mathfrak{X}[t_0, c_0] \backslash D, \Theta) = 0$$

or equivalently

$$H^0(\mathfrak{X}[t_0, c_0] \backslash D, \Omega^1) = 0.$$

Proof. Since we have a non-degenerate 2-form ω on $\mathfrak{X}[t_0, c_0]\backslash D$, the sheaf Θ is isomorphic to Ω^1. So we have to prove $H^0(\mathfrak{X}[t_0, c_0]\backslash D, \Omega^1) = 0$. Let $\eta \in H^0(\mathfrak{X}[t_0, c_0]\backslash D, \Omega^1)$. We show $\eta = 0$. In fact, $d\eta \in H^0(\mathfrak{X}[t_0, c_0]\backslash D, \Omega^2)$. It follows from Proposition (4.4) that we can find a complex number λ such that $d\eta = \lambda\omega$. By Corollary (5.25), $\lambda = 0$ or $d\eta = 0$. The algebraic surface $\mathfrak{X}[t_0, c_0]\backslash D$ is covered by three copies W_1, W_2, W_{12} of $\mathbb{A}^2$ so that we can find $\varphi_i \in H^0(W_i, \mathcal{O})$ for $i = 1, 3, 12$ such that $d\varphi_i = \eta$ on W_i for $i = 1, 3, 12$. Hence $\{\varphi_i - \varphi_j\}_{W_i \cap W_j}$ is a 1-cocycle or is an element of $Z^1(\bigcup W_i, \mathbb{C})$. Since $H^1_{zar}(\mathfrak{X}[t_0, c_0]\backslash D, \mathbb{C}) = 0$, we can find constants $k_i \in \mathbb{C}$ for $i = 1, 3, 12$ such that

$$\varphi_i - \varphi_j = k_i - k_j.$$

In other words the functions $\varphi_i - k_i$ glue together to give an element of $H^0(\mathfrak{X}[t_0, c_0]\backslash D, \mathcal{O})$. Now it follows from Corollary (4.8) that this function is constant k. Therefore we have $\varphi_i - k_i = k$ on W_i. This shows $\varphi_i = k + k_i \in \mathbb{C}$. Hence $0 = d\varphi_i = \eta$. This is what we had to show.

§6. Hamiltonian system.

We worked in §5 on $\mathfrak{X}[t_0, c_0]$ for a fixed t_0 and c_0. We can apply this argument to the relative case $\mathfrak{X}/\mathrm{Spec}\ \mathbb{C}[t, c]$. In theory of Painlevé equations, however we have to study $\mathfrak{X}/\mathrm{Spec}\ \mathbb{C}[c]$. We have three copies of $\mathbb{C}^4$:

$$\mathcal{W}_1 := \mathrm{Spec}\ \mathbb{C}[y_1, z_1, t, c],$$

$$\mathcal{W}_3 := \mathrm{Spec}\ \mathbb{C}[y_3, z_3, t, c],$$

$$\mathcal{W}_{12} := \mathrm{Spec}\ \mathbb{C}[y_{12}, z_{12}, t, c].$$

A point $(y_1, z_1, t_1, c_1) \in \mathcal{W}_1$ and a point $(y_3, z_3, t_3, c_3) \in \mathcal{W}_3$ and identified if $(t_1, c_1) = (t_3, c_3)$, $y_1 y_3 = 1$ and if $z_1 = y_3(c - y_3 z_3)$. A point (y_1, z_1, t_1, c_1) of $\mathcal{W}_1$ and a point $(y_{12}, z_{12}, t_{12}, c_{12})$ of $\mathcal{W}_{12}$ are identified if $(t_1, c_1) = (t_{12}, c_{12})$, $y_1 y_{12} = 1$ and if

$$z_1 + 2y_1^2 + t_1 = y_{12}\left(-(c+1) - y_{12} z_{12}\right).$$

A point (y_3, z_3, t_3, c_3) of $\mathcal{W}_3$ and a point $(y_{12}, z_{12}, t_{12}, c_{12})$ of $\mathcal{W}_{12}$ are identified if $(t_3, c_3) = (t_{12}, c_{12})$ and if

$$z_3 = z_{12} + \frac{2c_{12} + 1}{y_{12}} + \frac{t_{12}}{y_{12}^2} + \frac{2}{y_{12}^4}.$$

By gluing $\mathcal{W}_1$, $\mathcal{W}_2$ and $\mathcal{W}_3$ together by this rule, we have $\mathcal{W}_1 \cup \mathcal{W}_2 \cup \mathcal{W}_3 = \mathfrak{X}\backslash\mathfrak{D}$. Now we consider $\mathfrak{X}$ as a variety over $\mathrm{Spec}\ \mathbb{C}[c]$. In other words $\mathfrak{X} \to \mathrm{Spec}\ \mathbb{C}[c]$ is a family of threefolds parameterized by $\mathrm{Spec}\ \mathbb{C}[c]$. The differential, as well the $\wedge$, is take over $\mathbb{C}[c]$. To distinguish the differential over $\mathbb{C}[c]$ from the one over $\mathbb{C}[t,c]$, we denote the former by $d_{/c}$ so that $d_{/c}\mathbb{C}[c]=0$ but $d_{/c}t \neq 0$. Similarly we denote by $\wedge_{/c}$ the wedge over $\mathbb{C}[c]$.

We look for polynomials $H_1(t,c,y_1,z_1) \in \mathbb{C}[t,c,y_1,z_1]$, $H_3(t,c,y_3,z_3) \in \mathbb{C}[t,c,y_3,z_3]$, $H_{12}(t,c,y_{12},z_{12}) \in \mathbb{C}[t,c,y_{12},z_{12}]$ such that

$$d_{/c}y_1 \wedge_{/c} d_{/c}z_1 + d_{/c}H_1 \wedge_{/c} d_{/c}t, \tag{6.1}$$

$$d_{/c}y_3 \wedge_{/c} d_{/c}z_3 + d_{/c}H_3 \wedge_{/c} d_{/c}t$$

and

$$d_{/c}y_{12} \wedge_{/c} d_{/c}z_{12} + d_{/c}H_{12} \wedge_{/c} d_{/c}t$$

glue together and define a 2-form on $\mathfrak{X}\backslash\mathfrak{D}$ over Spec $\mathbb{C}[c]$.

Lemma (6.2). *Let* $H_1 = y_1^2 z_1 + \frac{1}{2}z_1^2 + \frac{t}{2}z_1 - cy_1 \in \mathbb{C}[t,c,y_1,z_1]$. *Then,*
(i) $H_3(t,c,y_3,z_3) := H_1(t,c,\frac{1}{y_3}, y_3(c-y_3z_3))$ *is a polynomial in* t, c, y_3, z_3.
(ii) $H_{12}(t,c,y_{12},z_{12}) := H_3\left(t,c,y_{12},z_{12}+\frac{2c+1}{y_{12}}+\frac{t}{y_{12}^2}+\frac{2}{y_{12}^4}\right) - \frac{1}{y_{12}}$ *is a polynomial in* t, c, y_{12}, z_{12}.

Proof. The lemma is proved by a simple calculation.

Lemma (6.3). *We have*

$$d_{/c}y_1 \wedge_{/c} d_{/c}z_1 = d_{/c}y_3 \wedge_{/c} d_{/c}z_3 \tag{6.3.1}$$

and

$$d_{/c}y_3 \wedge_{/c} d_{/c}z_3 = d_{/c}y_{12} \wedge_{/c} d_{/c}z_{12} - \frac{1}{y_{12}^2}d_{/c}y_{12} \wedge_{/c} d_{/c}t. \tag{6.3.2}$$

Proof. These formulas are consequences of the identification rule of $\mathcal{W}_1$, $\mathcal{W}_2$ and $\mathcal{W}_{12}$.

Proposition (6.4). *Closed 2-forms* $d_{/c}y_1 \wedge_{/c} d_{/c}z_1 + d_{/c}H_1 \wedge_{/c} d_{/c}t$, $d_{/c}y_3 \wedge_{/c} d_{/c}z_3 + d_{/c}H_3 \wedge_{/c} d_{/c}t$, $d_{/c}y_{12} \wedge_{/c} d_{/c}z_{12} + d_{/c}H_{12} \wedge_{/c} d_{/c}t$ *glue together and define a 2-form* ω_c *on* $\mathfrak{X}\backslash\mathfrak{D}$ *over* Spec $\mathbb{C}[c]$.

Proof. We have to show

$$d_{/c}y_1 \wedge_{/c} d_{/c}z_1 + d_{/c}H_1 \wedge_{/c} d_{/c}t = d_{/c}y_3 \wedge_{/c} d_{/c}z_3 + d_{/c}H_3 \wedge_{/c} d_{/c}t \tag{6.5.1}$$

on $\mathcal{W}_1 \cap \mathcal{W}_3$ and

$$d_{/c}y_3 \wedge_{/c} d_{/c}z_3 + d_{/c}H_3 \wedge_{/c} d_{/c}t = d_{/c}y_{12} \wedge_{/c} d_{/c}z_{12} + d_{/c}H_{12} \wedge_{/c} d_{/c}t \tag{6.5.2}$$

on $\mathcal{W}_3 \cap \mathcal{W}_{12}$. (6.5.1) follows from (6.3.1) and the fact that $H_1 = H_3$ (cf. Lemma (6.2), (ii)). On the other hand (6.5.2) follows from Lemma (6.2), (ii) and (6.3.2).

The 2-form ω_c defines a skew symmetric form on $\Theta_{\mathfrak{X}\backslash\mathfrak{D}/\mathrm{Spec}\ \mathbb{C}[c]}$, of which the null foliation is the second Painlevé equation. Namely $\Theta_{\mathfrak{X}\backslash\mathfrak{D}/\mathrm{Spec}\ \mathbb{C}[c]}$ is a vector bundle of rank three on which we have the skew symmetric form ω_c. So at each point p of $\mathfrak{X}\backslash\mathfrak{D}$, there exist a 1-dimensional subspace Σ_p of $\Theta_{\mathfrak{X}\backslash\mathfrak{D}/\mathrm{Spec}\ \mathbb{C}[t]}$ such that $\omega_c(\Sigma_p, \Theta_{\mathfrak{X}\backslash\mathfrak{D}/\mathrm{Spec}\ \mathbb{C}}, p) = 0$. The subspace Σ_p defines a foliation. Rigorously speaking, we have to fix $c = c_0 \in \mathbb{C}$ and work on the fiber $\mathfrak{X}_{c_0}\backslash\mathfrak{D}_{c_0}$, which is a threefold, over $c = c_0$ of $\mathfrak{X}\backslash\mathfrak{D} \to \mathrm{Spec}\ \mathbb{C}[c]$. Locally on $\mathcal{W}_1$ over $\mathbb{C}[c]$ for example the null foliation is given by the Hamiltonian flow

$$\frac{y_1}{dt} = \frac{\partial H_1}{\partial z_1}, \quad \frac{z_1}{dt} = -\frac{\partial H_1}{\partial y_1}. \tag{6.6}$$

Namely

$$\begin{cases} \dfrac{dy_1}{dt} = y_1^2 + z_1 + \dfrac{t}{2}, \\ \\ \dfrac{dz_1}{dt} = -2y_1 z_1 + c. \end{cases}$$

Theorem (6.7). *The Hamiltonian functions H_1, H_3, H_{12} are unique. Namely if H_1', H_3', H_{12}' are polynomials such that the 2-forms in (6.1) glue together, then $H_i - H_i'$ does not depend on y_i, z_i for $i = 1, 3, 12$.*

Proof. Let $H_i' \in \mathbb{C}[t, c, y_1, z_1]$, $H_3' \in \mathbb{C}[t, c, y_3, z_3]$, $H_{12}' \in \mathbb{C}[t, c, y_{12}, z_{12}]$ such that the 2-forms of (6.1) glue together. We have to show that $H_1 - H_1'$, $H_3 - H_3'$, $H_{12} - H_{12}'$ are functions of t, c. It follows from (6.1), we have $d(H_1 - H_1') = d(H_3 - H_3')$ on $\mathcal{W}_1 \cap \mathcal{W}_3$, $d(H_1 - H_1') = d(H_{12} - H_{12}')$ on $\mathcal{W}_1 \cap \mathcal{W}_{12}$, where d is taken over $\mathbb{C}[t, c]$ so that $d(\mathbb{C}[t, c]) = 0$. We set

$$\xi_{ij} = (H_i - H_i') - (H_j - H_j')$$

for $i, j \in \{1, 3, 12\}$. Then $\xi_{i,j} \in \mathbb{C}$ and $\{\xi_{ij}\}$ is a 1-cocycle with coefficients in $\mathbb{C}$. (Precisely speaking ξ_{ij} depends on t and c.) Since $H^1_{zar}(\mathfrak{X}[t, c], \mathbb{C}) = 0$, we can find constants $\xi_i \in \mathbb{C}$ such that $\xi_i - \xi_j = \xi_{ij}$ for $i, j \in \{1, 3, 12\}$. Hence $H_i - H_i' + \xi_i$ glue together and define a regular function on $\mathfrak{X}[t, c]\backslash D$. So $H_i - H_i' + \xi_i$ is a constant in the sense that it is a function of t and c by Corollary(4.8) for $i = 1, 3, 12$.

§7. Deformation of the open surface $\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0]$ and the Hamiltonian.

Let us fix $c = c_0$. Then we have a family $\mathfrak{X}_{c_0}\backslash\mathfrak{D}_{c_0}$ of open surfaces $\mathfrak{X}[t, c_0]\backslash\mathfrak{D}[t, c_0]$ parameterized by t. Namely we have

$$\mathfrak{X}_{c_0}\backslash\mathfrak{D}_{c_0} \to \mathrm{Spec}\ \mathbb{C}[t]. \tag{7.1}$$

Since we have a flow of the second Painlevé equation on $\mathfrak{X}_{c_0}\backslash\mathfrak{D}_{c_0}$ so that analytically the fibration (7.1) is trivial. If we take a point $t = t_0$, then

$$(\mathfrak{X}_{c_0}\backslash\mathfrak{D}_{c_0})^{an} \simeq (\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0])^{an} \times \mathbb{C}.$$

Now we consider the Kodaira-Spencer map

$$\varphi : T_{t_0} \to H^1\left(\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0], \Theta\right)$$

associated with the fibration (7.1). Here T_{t_0} denotes the tangent space of Spec $\mathbb{C}[t]$ at $t = t_0$. The open surface $\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0]$ is defined by gluing together three copies W_1, W_3, W_{12} of $\mathbb{C}^2$ by the rule in Theorem (4.22). We look at the Kodaira-Spencer map. The image $\varphi\left(\left(\frac{d}{dt}\right)_{t=t_0}\right) \in H^1\left(\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0], \Theta\right)$ is given by definition by the following 1-cocycle $\{\theta_{j,k}\} \in \Gamma(W_j \cap W_k, \Theta)$ $(j, k = 1, 3, 12)$. Let $y_j = f_{jk}(y_k, z_k, t)$, $z_j = g_{jk}(y_k, z_k, t)$. Then

$$\theta_{j,k} = \left.\frac{\partial f_{jk}}{\partial t}\frac{\partial}{\partial y_j} + \frac{\partial g_{jk}}{\partial t}\frac{\partial}{\partial z_j}\right|_{t=t_0} \in \Gamma(W_j \cap W_k, \Theta).$$

In our case, we have

$$0 = \theta_{1,3} = -\theta_{3,1} \in \Gamma(W_1 \cap W_3, \Theta).$$

$$\theta_{3,12} = \frac{1}{y_{12}^2}\frac{\partial}{\partial z_3} \in \Gamma(W_3 \cap W_{12}, \Theta), \qquad -\theta_{12,3} = \theta_{3,12}$$

and

$$0 = \theta_{1,12} = -\theta_{12,1} \in \Gamma(W_1 \cap W_{12}, \Theta).$$

The 1-cocycle $\{\theta_{i,j}\}$ is cohomologous to 0. Since we have a symplectic structure on $\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0]$, we have on $\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0]$ an isomorphism

$$\Theta \simeq \Omega^1. \tag{7.2}$$

The 1-cocycle $\{\omega_{ij}\}$ with coefficients in Ω^1 corresponding $\{\theta_{i,j}\}$ by isomorphism (7.2) is

$$\omega_{1,3} = -\omega_{3,1} = 0, \qquad \omega_{1,12} = -\omega_{12,1} = 0$$

$$\omega_{3,12} = \frac{1}{y_{12}^2}dy_{12} = \frac{1}{y_3^2}dy_3 \in H^0(W_3 \cap W_{12}, \Omega^1).$$

If we consider $\omega_j = dH_j$ on W_j $(j = 1, 3, 12)$, then $\omega_j - \omega_k = \omega_{ik}$ by Lemma (6.2). So the cohomology class determined by $\{\omega_{ij}\} \in H^1\left(\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0], \Omega^1\right)$ is 0 and hence the Kodaira-Spencer class $\{\theta_{ij}\} \in H^1\left(\mathfrak{X}[t_0, c_0]\backslash\mathfrak{D}[t_0, c_0], \Theta\right)$ is 0. So we can express this fact by saying that the Hamiltonian functions H_j trivialize the Kodaira-Spencer class of the family (7.1).

§8. Cohomology groups.

We fix t_0, c_0 and denote the rational surface $\mathfrak{X}[t_0, c_0]$ by X and $\mathfrak{D}[t_0, c_0]$ by D.

Lemma (8.1).

$$\chi(\Theta_X) = -10.$$

Proof. Since X is a rational surface, $\chi(\mathcal{O}_X) = 1$. For a vector bundle E of rank 2 over X, the Riemann-Roch theorem tells us

$$\chi(E) = \frac{1}{2}\left(c_1(E)^2 - 2c_2(E)\right) + \frac{1}{2}c_1(E)c_1(\Theta_X) + 2\chi(\mathcal{O}_X),$$

and hence

$$\chi(E) = \frac{1}{2}\left(c_1(E)^2 - 2c_2(E)\right) + \frac{1}{2}c_1(E)c_1(\Theta_X) + 2.$$

If we take Θ_X as E,

$$\chi(\Theta_X) = \frac{1}{2}\left(c_1(\Theta_X)^2 - 2c_2(\Theta_X)\right) + \frac{1}{2}c_1(\Theta_X)c_1(\Theta_X) + 2$$

so that

$$\chi(\Theta_X) = \frac{1}{2}\left(c_1(\Theta_X)^2 - 2c_2(\Theta_X)\right) + \frac{1}{2}c_1(\Theta_X)^2 + 2.$$

Since $c_1(\Theta_X) = -K$, $c_1(\Theta_X)^2 = 0$ and hence

$$\chi(\Theta_X) = -c_2(\Theta_X) + 2. \tag{8.2}$$

It follows from Noether's formula

$$\begin{aligned} 1 - q(X) + p_q(X) &= \frac{1}{12}\left(c_1(\Theta_X)^2 + c_2(\Theta_X)\right) \\ 12 &= c_2(\Theta_X). \qquad (8.3) \end{aligned}$$

Now the lemma follows from (8.2) and (8.3).

Proposition (8.4).

$$h^1(X, \Theta_X(-\log D)) = 2.$$

Proof. By the Serre duality

$$h^1\left(X, \Theta_X(-logD)\right) = h^1\left(X, K \otimes \Omega^1_X(\log D)\right)$$

and hence we have to calculate $h^1\left(X, K \otimes \Omega^1_X(\log D)\right)$. We have an exact sequence

$$0 \to \Omega^1_X \to \Omega^1_X(\log D) \to \bigoplus_{i=1}^{8} \mathcal{O}_{D_i} \to 0.$$

Tensoring K with the exact sequence, we get

$$0 \to K \otimes \Omega^1_X \to K \otimes \Omega^1_X(\log D) \to \bigoplus_{i=1}^{8} K \otimes \mathcal{O}_{D_i} \to 0.$$

Since D_i is a -2-curve, we have $K \otimes \mathcal{O}_{D_i} \simeq \mathcal{O}_{D_i}$ and consequently we have

$$0 \to K \otimes \Omega^1_X \to K \otimes \Omega^1_X(\log D) \to \bigoplus_{i=1}^{8} \mathcal{O}_{D_i} \to 0. \tag{8.5.1}$$

This exact sequence gives the long exact sequence

$$\begin{aligned} 0 &\to H^0\left(K \otimes \Omega^1_X\right) \to H^0\left(K \otimes \Omega^1_X(\log D)\right) \to H^0\left(\bigoplus_{i=1}^{8} \mathcal{O}_{D_i}\right) \\ &\to H^1\left(K \otimes \Omega^1_X\right) \to H^1\left(K \otimes \Omega^1_X(\log D)\right) \to H^1\left(\bigoplus_{i=1}^{8} \mathcal{O}_{D_i}\right) \\ &\to H^2\left(K \otimes \Omega^1_X\right) \to H^2\left(K \otimes \Omega^1_X(\log D)\right) \to 0. \end{aligned} \tag{8.5.2}$$

Since $K = -\sum_{i=0}^{7} n_i D_i$ with $n_i > 0$ for $0 \le i \le 7$, $K \otimes \Omega^1_X(\log D)$ is a subsheaf of Ω^1_X so that $H^0\left(K \otimes \Omega^1_X(\log D)\right) = 0$. The D_i's are isomorphic to $\mathbb{P}^1$ and hence $H^0\left(\bigoplus_{i=1}^{8} \mathcal{O}_{D_i}\right) = \mathbb{C}^8$, $H^1\left(\bigoplus_{i=1}^{8} \mathcal{O}_{D_i}\right) = 0$. So it follows from the long exact sequence

$$0 \to \mathbb{C}^8 \to H^1\left(K \otimes \Omega^1_X\right) \to H^1\left(K \otimes \Omega^1_X(\log D)\right) \to 0.$$

We had to show $h^1\left(K \otimes \Omega^1_X(\log D)\right) = 2$. To this end, it suffices to show $h^1\left(K \otimes \Omega^1_X\right) = 10$ by the above exact sequence. It follows from the Serre duality $h^1\left(K \otimes \Omega^1_X\right) = h^1\left(\Theta_X\right)$ so that we have to show

$$h^1(\Theta_X) = 10.$$

We notice here by the Serre duality $h^2\left(\Theta_X\right) = h^0\left(K \otimes \Omega^1_X\right)$. Since $-K$ is effective $K \otimes \Omega^1_X$ is a subsheaf of Ω^1_X and since $H^0\left(\Omega^1_X\right) = 0$ because X is rational, $H^0\left(K \otimes \Omega^1_X\right) = 0$. Therefore

$$h^2\left(\Theta_X\right) = 0. \tag{8.6}$$

Now (8.4) follows from Theorem (5.26), Lemma (8.1) and (8.6).

Corollary (8.7). *to the proof.* $H^0\left(\Theta_X(-\log D)\right) = H^2\left(\Theta_X(-\log D)\right) = 0.$

Proof. In fact since $H^0(\Theta_X(-\log D)) \subset H^0\left(X\backslash D, \Theta_X\right)$, $H^0\left(\Theta_X(-\log D)\right) = 0$ by Theorem (5.26). We noticed in the proof of Proposition (8.4)

$$H^0\left(K \otimes \Omega^1_X(\log D)\right) = 0,$$

implies by the Serre duality $H^2\left(\Theta_X(-\log D)\right) = 0$.

Corollary (8.8). *to the proof. If* $c_0 = 0$, *we know that there is a* -2*-curve* C'_2 *on* $X[t_0, 0]\backslash D$. *We have*

$$h^1\left(X, \Theta_X\left(-\log(D + C'_2)\right)\right) = 1$$

and

$$h^0\left(\Theta_X\left(-\log(D + C'_2)\right)\right) = h^2\left(\Theta_X\left(-\log(D + C'_2)\right)\right) = 0.$$

Proof. We argue as in the proof of Proposition (8.4). We replace D by $D + C'_2$ and consider the exact sequence (8.5.1) and the long exact sequence (8.5.2). Then we have to show $H^0\left(K \otimes \Omega^1_X\left(\log(D + C'_2)\right)\right) = 0$. Since $K \otimes \Omega^1_X\left(\log(D + C'_2)\right)$ is a subsheaf of $\Omega^1_X(\log C'_2)$, it is sufficient to show $H^0\left(\Omega^1_X(\log C'_2)\right) = 0$. In fact we have a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & H^0(\Omega^1_X) \arrow[r] \arrow[d] & H^0\left(\Omega^1_X(\log C'_2)\right) \arrow[r] \arrow[d] & H^0\left(\mathcal{O}_{C'_2}\right) \arrow[r, phantom, "\simeq"] \arrow[d, equal] & \mathbb{C} \\
 & H^1(X, \mathbb{C}) \arrow[r] & H^1\left(X\backslash C'_2, \mathbb{C}\right) \arrow[r] & H^2\left(X; C'_2, \mathbb{C}\right) \arrow[r] & H^2(X, \mathbb{C})
\end{tikzcd}$$

of cohomology groups for the usual topology. By the Poincaré duality we have $H^2(X; C'_2, \mathbb{C}) \simeq H_2(C'_2, \mathbb{C})$ and hence the morphism $H^2(X; C'_2, \mathbb{C}) \simeq H_2(C'_2, \mathbb{C}) \simeq \mathbb{C} \to H^2(X, \mathbb{C})$ is injective. So the morphism $H^1(X\backslash C'_2, \mathbb{C}) \to H^2(X; C'_2, \mathbb{C})$ is trivial. Since $H^0(\Omega^1_X) = 0$, this shows $H^0(\Omega^1_X(\log C'_2)) = 0$.

§9. Rational singular points.

Let Y be an affine surface defined over $\mathbb{C}$ and P be a point. We assume that Y is normal, $Y\backslash\{D\}$ is smooth and that P is a rational double point of type A_1. Let $f : \widetilde{Y} \to Y$ be the minimal resolution. So $C := \varphi^{-1}(P)$ is a curve isomorphic to $\mathbb{P}^1$ with $C^2 = -2$. Ω^1_Y is the sheaf of Kähler differentials on Y over $\mathbb{C}$. Θ_Y is the dual of Ω^1_Y so that $\Theta_Y = \mathrm{Hom}_{\mathcal{O}_Y}(\Omega^1_Y, \mathcal{O}_Y)$.

Proposition (9.1). *We have*

$$f_*\Theta_{\widetilde{Y}}(-\log C) \simeq \Theta_Y$$

and

$$R^1 f_*\Theta_{\widetilde{Y}}(-\log C) = 0.$$

Proof. It follows form an exact sequence

$$0 \to \Theta_{\widetilde{Y}}(-\log C) \to \Theta_{\widetilde{Y}} \to N_C \to 0,$$

the long exact sequence

$$\begin{aligned} 0 &\to f_*\Theta_{\widetilde{Y}}(-\log C) \to f_*\Theta_{\widetilde{Y}} \to f_*N_C \\ &\to R^1f_*\Theta_{\widetilde{Y}}(-\log C) \to R^1f_*\Theta_{\widetilde{Y}} \xrightarrow{i} R^1f_*N_C \to 0, \end{aligned}$$

N_C being the normal bundle of the curve C. Since $N_C \simeq \mathcal{O}_C(-2)$, we have $f_*N_C = 0$. By Proposition(1.2) in [BW],

$$f_*\Theta_{\widetilde{Y}} \simeq \Theta_Y.$$

Moreover for the minimal resolution of a rational double point, we have an isomorphism $H^1(\Theta_{\widetilde{Y}}) \simeq H^1(N_C)$ by (1.8) in [BW]. Hence the morphism i in the long exact sequence above is an isomorphism and consequently

$$f_*\Theta_{\widetilde{Y}}(-\log C) \simeq f_*\Theta_{\widetilde{Y}} \simeq \Theta_Y, \qquad R^1f_*\Theta_{\widetilde{Y}}(-\log C) = 0.$$

§10. Calculation of Ext.

On rational surface $\mathfrak{X}[t_0,0]$, there is a curve C_2' isomorphic $\mathbb{P}^1$ with $(C_2')^2 = -2$ and $C_2' \cap D = \emptyset$. We can contract the curve C_2' on $\mathfrak{X}[t_0,0]$ to a rational singular point P to get a rational surface $\mathcal{Y}[t_0,0]$ with a rational double point P of type A_1. We have a morphism

$$f : \mathfrak{X}[t_0,0] \to \mathcal{Y}[t_0,0],$$

which is the minimal resolution of $\mathcal{Y}[t_0,0]$. Since the curve D does not intersect C_2', f is an isomorphism on a neighborhood of D. So we denote the image $f(D)$ on $\mathcal{Y}[t_0,0]$ again by D. We compare the Kuranishi family of the pair $(\mathfrak{X}[t_0,0], D)$ and that of the pair $(\mathcal{Y}[t_0,0], D)$. We denote $\mathfrak{X}[t_0,0]$ by X and $\mathcal{Y}[t_0,0]$ by Y.

Lemma (10.1).

$$H^2(Y, \Theta_Y(-\log D)) = 0$$

Proof. It follows from Proposition (9.1) that the spectral sequence

$$\begin{aligned} E_2^{p,q} &= H^p\left(Y, R^qf_*\Theta_X\left(-\log(D + C_2')\right)\right) \\ &\Rightarrow H^n\left(X, \Theta_X\left(-\log(D + C_2')\right)\right) \end{aligned}$$

degenerates and that $f_*\Theta_X\left(-\log(D + C_2')\right) = \Theta_Y(-\log D)$. So we have

$$H^p\left(Y, \Theta_Y(-\log D)\right) = H^p\left(X, \Theta_X\left(-\log(D + C_2')\right)\right)$$

for every integer $p \geq 0$. The lemma now follows from Corollary (8.8).

Proposition (10.2). *The spectral sequence*

$$E_2^{p,q} = H^p\left(Y, \mathcal{E}xt^q_{\mathcal{O}_Y}\left(\Omega^1_Y(\log D), \mathcal{O}_Y\right)\right)$$

$$\Rightarrow \mathrm{Ext}^{p+q}_{\mathcal{O}_Y}\left(\Omega^1_Y(\log D), \mathcal{O}_Y\right)$$

degenerates at the E_2-terms and we have the exact sequence

$$0 \to H^1\left(Y, \Omega^1_Y(\log D)\right) \to \mathrm{Ext}^1_{\mathcal{O}_Y}\left(\Omega^1(\log D), \mathcal{O}_Y\right) \to \mathrm{Ext}^1_{\mathcal{O}_{Y,P}}\left(\Omega^1_{Y,P}, \mathcal{O}_{Y,P}\right) \to 0.$$

Proof. Since Y is locally a complete intersection, the projective dimension of $\Omega^1_Y(-\log D)$ is equal to the projective dimension of $\Omega^1_Y \leq 1$ so that

$$\mathcal{E}xt^q_{\mathcal{O}_Y}\left(\Omega^1_Y(-\log D), \mathcal{O}_Y\right) = 0$$

for $q \geq 2$. $\mathcal{E}xt^1_{\mathcal{O}_Y}\left(\Omega^1_Y(-\log D), \mathcal{O}_Y\right)$ is supported on the singular point P. Lemma (10.1) shows $E_2^{0,2} = 0$. So the spectral sequence degenerates at the E_2-terms. Thus $\mathrm{Ext}^2_{\mathcal{O}_Y}\left(\Omega^1(\log D), \mathcal{O}_Y\right) = 0$ and we have an exact sequence

$$0 \to E_2^{1,0} \to \mathrm{Ext}^1_{\mathcal{O}_Y}\left(\Omega^1_Y(-\log D), \mathcal{O}_Y\right) \to E_2^{0,1} \to 0,$$

which is nothing but the exact sequence of the proposition.

§11. Family $\mathfrak{X}[t,c]$ of rational surfaces.

Theorem (11.1). *The family* $(\mathfrak{X}[t,c], \mathfrak{D}[t,c]) \to \mathrm{Spec}\ \mathbb{C}[t,c]$ *of pairs a surface and divisor is a Kuranishi family at every point* $(t_0, c_0) \in (\mathbb{C}[t,c])^{an} = \mathbb{C}^2$.

Proof. Again we denote $\mathfrak{D}[t_0, c_0]$ by D. The theorem says that the Kodaira-Spencer map

$$\rho : T_{\mathbb{C}^2,P} \to H^1\left(\mathfrak{X}[t_0, c_0], \Theta(-\log D)\right) \tag{11.2}$$

is an isomorphism of vector spaces, where $T_{\mathbb{C}^2,P}$ is the tangent space of $(\mathrm{Spec}\ \mathbb{C}[t,c])^{an} = \mathbb{C}^2$ at a point $P = (t_0, c_0)$. It follows from Proposition (8.4) that we have to show that the image of Kodaira-Spencer map is of dimension 2. As we have show in §2, [U2], the surface $\mathfrak{X}[t_0, c_0]$ is covered by 12 affine open sets W_i isomorphic to $\mathbb{A}^2$ $(1 \leq i \leq 12)$. The image

$$\rho\left(\left.\left(\frac{\partial}{\partial c}\right)\right|_{(t,c)=(t_0,c_0)}\right) \in H^1\left(\mathfrak{X}[t_0, c_0], \Theta(-\log D)\right)$$

is represented by a Čech 1-cocycle $a = \{a_{i,j}\}_{1 \leq i,j \leq 12}$ with coefficients in $\Theta(-\log D)$ with respect to the covering $X[t_0, c_0] = \cup_{i=1}^{12} W_i$ so that

$$a_{i,j} \in H^0\left(W_i \cap W_j, \Theta(-\log D)\right).$$

Similarly, the image

$$\rho\left(\left(\frac{\partial}{\partial t}\right)\bigg|_{(t,c)=(t_0,c_0)}\right) \in H^1\left(\mathfrak{X}[t_0,c_0],\Theta(-\log D)\right)$$

is given by Čech 1-cocycle

$$b = \{b_{i,j}\}_{1\le i,j\le 12}$$

with

$$b_{i,j} \in H^0\left(W_i \cap W_j, \Theta(-\log D)\right).$$

We assume that a linear combination $\lambda a + \mu b$, which is a Čech 1-cocycle with coefficients in $\Theta(-\log D)$, is cohomologous to 0 for complex numbers λ, μ and we show that $\lambda = \mu = 0$. We recall that $\mathfrak{X}[t_0,c_0]\backslash D$ is covered by W_1, W_3 and W_{12}. We have a canonical map

$$H^1\left(\mathfrak{X}[t_0,c_0],\Theta(-\log D)\right) \to H^1\left(\mathfrak{X}[t_0,c_0]\backslash D,\Theta(-\log D)\right)$$

The image of a Čech 1-cocycle $f = \{f_{i,j}\}$ under this map is denoted by $\bar{f} = \{\bar{f}_{i,j}\}$. Since as we have seen in §7, $\bar{b}$ is cohomologous to 0, it follows from the assumption that $\lambda\bar{a} = \lambda\bar{a} + \mu\bar{b}$ is cohomologous to 0. By the coordinate transformation between W_1 and W_3 given in §5, we get

$$\frac{\partial z_1}{\partial c} = \frac{\partial}{\partial c}(y_3 c - y_3 z_3) = y_3 = \frac{1}{y_1}$$

and $\partial y_1/\partial c = 0$. So by the definition of the Kodaira-Spencer map, we have

$$\bar{a}_{1,3} = \frac{1}{y_1}\frac{\partial}{\partial z_1} \in H^0\left(W_1 \cap W_3, \Theta(-\log D)\right). \tag{11.3}$$

Similarly

$$\bar{a}_{1,12} = -\frac{1}{y_1}\frac{\partial}{\partial z_1} \in H^0\left(W_1 \cap W_{12}, \Theta(-\log D)\right). \tag{11.4}$$

On the other hand, we have the symplectic structure on $\mathfrak{X}[t_0,c_0]\backslash D = W_1 \cup W_2 \cup W_3$ given by $dy_1 \wedge dz_1 = dy_3 \wedge dz_3 = dy_{12} \wedge dz_{12}$ so that we have an isomorphism

$$\varphi : \Theta \simeq \Omega^1 \tag{11.5}$$

on $\mathfrak{X}[t_0,c_0]\backslash D = W_1 \cup W_2 \cup W_3$. Under this isomorphism $\partial/\partial y_i$ corresponds to dz_i and $\partial/\partial z_i$ to $-dy_i$. We set $\bar{\alpha}_{i,j} := \varphi(\bar{a}_{i,j}) \in H^0(W_i \cap W_j, \Omega^1)$ for $i,j = 1,3,12$. In fact we have explicitly

$$\bar{\alpha}_{1,3} = -\frac{1}{y_1}dy_1, \qquad \bar{\alpha}_{1,12} = \frac{1}{y_1}dy_1. \tag{11.6}$$

Since $\lambda\bar{\alpha}$ is cohomologous to 0, there exist $\bar{\alpha}_i \in H^0(W_i, \Omega^1)$ for $i = 1, 3, 12$ such that

$$\bar{\alpha}_1 - \bar{\alpha}_3 = -\frac{\lambda}{y_1}dy_1, \qquad \bar{\alpha}_1 - \bar{\alpha}_{12} = \frac{\lambda}{y_1}dy_1. \tag{11.7}$$

In view of (11.6) and (11.7), $d\bar{\alpha}_1 = d\bar{\alpha}_3 = d\bar{\alpha}_{12}$ glue together and define a 2-form on $\mathfrak{X}[t_0, c_0]\backslash D$. So by Corollary (4.7), $d\bar{\alpha}_1 = d\bar{\alpha}_3 = d\bar{\alpha}_{12} = \nu\omega$ for a complex number ν. Since ω is not exact by Corollary (5.26), we have $\nu = 0$ and $d\bar{\alpha}_1 = d\bar{\alpha}_3 = d\bar{\alpha}_{12} = 0$. Now since $W_i \simeq \mathbb{A}^2$, by a theorem of Grothendieck, there exist polynomials $f_1 \in \mathbb{C}[y_1, z_1]$, $f_3 \in \mathbb{C}[y_3, z_3]$, $f_{12} \in \mathbb{C}[y_{12}, z_{12}]$ such that $df_i = \bar{\alpha}_i$ for $i = 1, 3, 12$. So it follows from (11.7)

$$df_1 - df_2 = -\frac{\lambda}{y_1}dy_1$$

on $W_1 \cap W_3 = \operatorname{Spec} \mathbb{C}[y_1, 1/y_1, z_1]$. Since $-\frac{\lambda}{y_1}dy_1$ is exact on $W_1 \cap W_3$ if and only if $\lambda = 0$. We conclude $\lambda = 0$. Now it remains to show $\mu = 0$. Let us consider the canonical map

$$H^1\left(\mathfrak{X}[t_0, c_0], \Theta(-\log D)\right) \to H^1\left(W_1 \cup W_2 \cup W_3 \cup W_{12}, \Theta(-\log D)\right).$$

The image of a 1-Čech cocycle $f = \{f_{i,j}\}$ under this map will be denoted by $\widetilde{f} = \{\widetilde{f_{i,j}}\}$. We calculated $\widetilde{b}$ for $i = 1, 3, 12$ in §6. We recall the coordinate transformation between W_1 and W_2, which are a part of coverings of the starting ruled surface in the construction of $X[t_0, c_0]$, is given by $y_1 = y_2$ and $z_1 = 1/z_2$ (cf. §1). It follows from the definition of the Kodaira-Spencer map $\widetilde{b}_{1,2} = 0$. Since μb is cohomologous to 0, $\mu\widetilde{b}$ is cohomologous to 0 too. We can find $\widetilde{b}_i \in H^0\left(W_i, \Theta(-\log D)\right)$ for $i = 1, 2, 3, 12$ such that

$$\widetilde{b}_i - \widetilde{b}_j = \mu\widetilde{b}_{i,j} \tag{11.8}$$

for $i, j = 1, 2, 3, 12$. Since $H^0\left(\mathfrak{X}[t_0, c_0]\backslash D, \Theta\right) = H^0\left(\mathfrak{X}[t_0, c_0]\backslash D, \Theta(-\log D)\right) = 0$, $\widetilde{b}_i \in H^0\left(W_i, \Theta\right)$ is uniquely determined for $i = 1, 3, 12$. Namely we have

$$\widetilde{b}_{t,i} = \mu\left(\frac{\partial H_i}{\partial z_i}\frac{\partial}{\partial y_i} - \frac{\partial H_i}{\partial y_i}\frac{\partial}{\partial z_i}\right)$$

for $i = 1, 3, 12$, where H_i is the Hamiltonian as we have seen in §7. In particular

$$\widetilde{b}_1 = \mu\left\{\left(y_1^2 + z_1 + \frac{t}{2}\right)\frac{\partial}{\partial y_1} + (c - 2y_1z_1)\frac{\partial}{\partial z_1}\right\}.$$

Since as we noticed above $\widetilde{b}_{1,2} = 0$, we have

$$\widetilde{b}_1 - \widetilde{b}_2 = 0,$$

which means $\widetilde{b}$ is regular on W_2. We have

$$\widetilde{b}_1 = \mu\left\{\left(y_2^2 + \frac{1}{z_2} + \frac{t}{2}\right)\frac{\partial}{\partial y_2} + \left(c - 2y_2\frac{1}{z_2}\right)\left(-z_2^2\right)\frac{\partial}{\partial z_2}\right\}$$

on W_2. So if $\widetilde{b}_1$ is regular on W_2, then $\mu = 0$. This is what we had to prove.

It is an interesting problem to give a pair of algebraic varieties $(U,\, V)$ defined over $\mathbb{C}$ such that the algebraic varieties U and V are not isomorphic one another but the associated analytic spaces U^{an} and V^{an} are isomorphic. A well-known example due to Serre is related with algebraic groups(See [H], Chapter VI, §3). The family $(\mathfrak{X}[t, c],\, \mathfrak{D}[t,\, c])$ provides us many such examples. Let us choose a point $(t_0,\, c_0) \in \mathbb{C}^2$. Let $t_1 \neq t_0$ be an arbitrary point in a small neighbourhood of $t_0 \in \mathbb{C}$. The pair $\mathfrak{X}[t_0,\, c_0] \setminus \mathfrak{D}[t_0,\, c_0],\, \mathfrak{X}[t_1,\, c_0] \setminus \mathfrak{D}[t_1,\, c_0])$ is a such pair. In fact, $\mathfrak{X}[t_0,\, c_0] \setminus \mathfrak{D}[t_0,\, c_0]$ is not isomorphic to $\mathfrak{X}[t_1,\, c_0] \setminus \mathfrak{D}[t_1,\, c_0]$ as algebraic varieties. In fact if we had an isomorphism

$$\varphi : \mathfrak{X}[t_0,\, c_0] \setminus \mathfrak{D}[t_0,\, c_0] \to \mathfrak{X}[t_1,\, c_0] \setminus \mathfrak{D}[t_1,\, c_0]$$

as algebraic varieties, then since the complementary divisors $\mathfrak{D}[t_0,\, c_0]$ and $\mathfrak{D}[t_1,\, c_0]$ consist of -2-curves, we could extend φ to an isomorphism

$$\bar{\varphi} : \mathfrak{X}[t_0,\, c_0] \to \mathfrak{X}[t_1,\, c_0]$$

of algebraic varieties so that the pair $(\mathfrak{X}[t_0,\, c_0],\, \mathfrak{D}[t_0,\, c_0])$ is isomorphic to the pair $(\mathfrak{X}[t_1,\, c_0],\, \mathfrak{D}[t_1,\, c_0])$. This contradicts Theorem(11.1). Integration of the second Painlevé equation that has no movable singular points gives an analytic isomorphism

$$\mathfrak{X}[t_0,\, c_0] \setminus \mathfrak{D}[t_0,\, c_0] \simeq \mathfrak{X}[t_1,\, c_0] \setminus \mathfrak{D}[t_1,\, c_0].$$

References

[A] D. Arinkin and S. Lysenko, *Isomorphisms between moduli spaces of $SL(2)$-bundles with connections on $\mathbb{P}^1 \backslash \{x_1, \ldots, x_q\}$*, Math. Res. Letters **4**, (1997), 181–190.

[BW] D.M. Burns and J.M. Wahl, *Local contributions to global deformations of surfaces* Invent. Math. 26(1974), 67-88.

[H] R. Hartshorne, *Ample subvarieties of algebraic varieties* Springer LNM 156, Springer-Verlag, Berlin-Heidelberg-NewYork, 1970.

[K] J. Kollar, *The structure of algebraic threefolds,* Bull.Aner.Math.Soc. **17**, (1987), 210-273.

[MMT] T. Matano, A. Matumiya and K. Takano, *On some Hamiltonian structures of Painlevé systems,* Funkcial. Ekvac., **40**, (1997) 271-291.

[O] K. Okamoto, *Sur les feuilleges associés aux équations du second ordre à points critiques fixes de P. Painlevé,* Japan. J. Math. 5, (1979), 1-79.

[STT] M.-H. Saito, T. Takebe, & H. Terajima, *Deformation of Okamoto–Painlevé pairs and Painlevé equations*, preprint, Kobe 2000, math.AG 0006026

[Sakai] H. Sakai, Rational surfaces associated with affine root systems and geometry of the Painlevé equations, preprint, Kyoto-Math 99-10.

[ST] T. Shioda and K, Takano, *On some Hamiltonian structures of Painlevé systems*, J. Math. Soc. Japan, **51**, (1999), no. 4, 843–866

[U1] H. Umemura, *On the irreducibility of the first Painlevé equation,* Algebraic Geometry and Commutative Algebra in honor of Masayoshi NAGATA, Tokyo : Konokuniya 1987, 771-789.

[U2] —-, *On the transformation group of the second Painlevé equation,* Nagoya Math. J. 157(2000), 15-46.

[UW] H. Umemura and H. Watanabe, *Solution of the second and fourth Painlevé equations, I,* Nagoya Math. J. 148 (1997), 151-198.